

DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

Spatial Besov Regularity for Semilinear Stochastic Partial Differential Equations on Bounded Lipschitz Domains

P. A. Cioica, S. Dahlke

Preprint 99



Edited by

AG Numerik/Optimierung
Fachbereich 12 - Mathematik und Informatik
Philipps-Universität Marburg
Hans-Meerwein-Str.
35032 Marburg


DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

Spatial Besov Regularity for Semilinear Stochastic Partial Differential Equations on Bounded Lipschitz Domains

P. A. Cioica, S. Dahlke

Preprint 99



The consecutive numbering of the publications is determined by their chronological order.

The aim of this preprint series is to make new research rapidly available for scientific discussion. Therefore, the responsibility for the contents is solely due to the authors. The publications will be distributed by the authors.

Spatial Besov Regularity for Semilinear Stochastic Partial Differential Equations on Bounded Lipschitz Domains

PETRU A. CIOICA, STEPHAN DAHLKE

Abstract

We study the spatial regularity of semilinear parabolic stochastic partial differential equations on bounded Lipschitz domains $\mathcal{O} \subseteq \mathbb{R}^d$ in the scale $B_{\tau,\tau}^\alpha(\mathcal{O})$, $1/\tau = \alpha/d + 1/p$, $p \geq 2$ fixed. The Besov smoothness in this scale determines the order of convergence that can be achieved by adaptive numerical algorithms and other nonlinear approximation schemes. The proofs are performed by establishing weighted Sobolev estimates and combining them with wavelet characterizations of Besov spaces.

Keywords: Semilinear stochastic partial differential equation, Besov space, weighted Sobolev space, nonlinear approximation, wavelet expansions

Mathematics Subject Classification (2010): 60H15, *Secondary:* 46E35, 65C30

1 Introduction

This paper is concerned with the spatial regularity of the solutions to semilinear stochastic partial differential equations (semilinear SPDEs, for short) measured in specific scales of Besov spaces. The motivation for these studies can be explained as follows.

In recent years, the numerical treatment of operator equations by adaptive numerical algorithms has become a field of increasing importance, with many applications in science and engineering. Especially, adaptive finite element schemes have been very successfully developed and implemented, and innumerable numerical experiments impressively confirm their excellent performance. Complementary to this, also adaptive algorithms based on wavelets have become more and more in the center of attention during the last years, for

This work has been supported by the Deutsche Forschungsgemeinschaft (DFG, grants DA 360/13-1, RI 599/4-1, SCHI 419/5-1) and a doctoral scholarship of the Philipps-Universität Marburg.

the following reason. The strong analytical properties of wavelets can be used to derive adaptive strategies which are guaranteed to converge for a huge class of elliptic operator equations, involving operators of negative order [4, 11]. Moreover, these algorithms are optimal in the sense that they asymptotically realize the convergence order of the optimal (but not directly implementable) approximation scheme, i.e. the order of best N -term wavelet approximation. Moreover, the number of arithmetic operations that is needed stays proportional to the number of degrees of freedom [4]. By now, various generalizations to non-elliptic equations [5], saddle point problems [12] and also nonlinear operator equations [6] exist. For finite element schemes, rigorous statements of these forms have been rather rare, even though, inspired by the results for wavelet schemes, the situation has changed during the last years [1, 18].

Although the above mentioned results are quite impressive, in the realm of adaptivity one is always faced with the following question: does adaptivity really pay for the problem under consideration, i.e. does our favourite adaptive scheme really provide a substantial gain of efficiency compared to more conventional nonadaptive schemes which are usually much easier to implement? At least for the case of adaptive wavelet schemes, it is possible to give a quite rigorous answer. A reasonable comparison would be to compare the performance of wavelet algorithms with classical, nonadaptive schemes which consist of approximations by linear spaces that are generated by uniform grid refinements. It is well-known that, under natural assumptions, the approximation order that can be achieved by such a uniform method depends on the smoothness of the exact solution measured in the classical L_p -Sobolev scale. More precisely, let $e_N(f)$ denote the uniform approximation error, then

$$f \in W_p^s(\mathcal{O}) \implies e_N(f) \leq C \cdot N^{-s/d}.$$

On the other hand, as already outlined above, for adaptive wavelet methods the best N -term approximation serves as the benchmark scheme. It is well-known that the convergence order that can be achieved by best N -term approximations also depends on the smoothness of the object we want to approximate. But, in contrast to the case of non-adaptive schemes, the smoothness has now to be measured in specific Besov spaces, usually corresponding to L_τ -spaces with $0 < \tau < 2$. Indeed, let $\sigma_N(f)$ denote the error of best N -term approximation, then

$$f \in B_{\tau,\tau}^\alpha(\mathcal{O}), \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p} \implies \sigma_N(f) \leq C \cdot N^{-\alpha/d}.$$

For a detailed description of these relationships, we refer, e.g. to [10] and [16], see also [3, Chapters 3 and 4]. Consequently, we can make the following statement: the use of adaptive wavelet schemes is completely justified if the spatial Besov smoothness α of the unknown solution of our SPDE in the scale $\frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}$ is higher compared to its regularity in the Sobolev scale. The situation is illustrated in Figure 1, where each point $(1/\tau, s)$ represents the smoothness spaces of functions with “ s derivatives in $L_\tau(\mathcal{O})$ ”.

For deterministic elliptic partial differential equations, many positive results in this direction already exist, see e.g. [7], [8], [9], [13], [14]. In contrast, little is known concerning




Figure 1: Linear vs. nonlinear approximation illustrated in a DeVore-Triebel diagram.

SPDEs. To our knowledge, the first results were obtained in [2]. In that paper, linear stochastic partial differential equations are considered and sufficiently high Besov smoothness to justify the use of adaptive schemes is established. The aim of this paper is to generalize these basic results to *semilinear* equations.

The stochastic setting is as follows. Let $\mathcal{O} \subset \mathbb{R}^d$ be a bounded Lipschitz domain. Fix $T \in (0, \infty)$ and let $(\mathcal{F}_t)_{t \in [0, T]}$ be a normal filtration on a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Furthermore, let $(w_t^k)_{t \in [0, T]}$, $k \in \mathbb{N}$, be a sequence of independent \mathbb{R} -valued standard Brownian motions w.r.t. $(\mathcal{F}_t)_{t \in [0, T]}$. For arbitrary $\varepsilon = (\varepsilon_f, \varepsilon_g) \in [0, \infty) \times [0, \infty)$ we consider the model equation

$$du = (Au + \varepsilon_f F(u))dt + \sum_{k=1}^{\infty} (\varepsilon_g G(u)^k + g^k(t))dw_t^k, \quad u(0, \cdot) = u_0, \quad (*_{\varepsilon})$$

for $t \in [0, T]$ on the domain \mathcal{O} . Here du is Itô's stochastic differential with respect to t , A denotes the Laplacian and the coefficients g^k , $k \in \mathbb{N}$, are random functions depending on $t \in [0, T]$ and $x \in \mathcal{O}$. The functions F and G are Lipschitz continuous on certain classes of stochastic processes taking values in weighted Sobolev spaces. For details see Section 2, especially Assumptions 5.

Equation $(*_{\varepsilon})$ is understood in a weak or distributional sense, i.e. u_{ε}^* is a solution of $(*_{\varepsilon})$, if for all smooth and compactly supported test functions $\varphi \in \mathcal{C}_0^{\infty}(\mathcal{O})$ the equality

$$\langle u_{\varepsilon}^*(t, \cdot), \varphi \rangle = \langle u_0, \varphi \rangle + \int_0^t \langle \Delta u_{\varepsilon}^*(s, \cdot) + \varepsilon_f F(u_{\varepsilon}^*)(s, \cdot), \varphi \rangle ds$$

$$+ \sum_{k=1}^{\infty} \int_0^t \langle \varepsilon_g G(u_\varepsilon^*)^k(s, \cdot) + g^k(s, \cdot), \varphi \rangle dw_s^k$$

holds for all $t \in [0, T]$ \mathbb{P} -almost surely. Here and throughout the paper we write $\langle u, \varphi \rangle$ for the application of a distribution $u \in \mathcal{D}'(\mathcal{O})$ to a test function $\varphi \in \mathcal{C}_0^\infty(\mathcal{O})$. We refer again to Section 2 for details.

Let us assume that the solution $u_\varepsilon^* = u_\varepsilon^*(t, \omega, x)$, $(t, \omega, x) \in [0, T] \times \Omega \times \mathcal{O}$, vanishes on the boundary $\partial\mathcal{O}$, satisfying a zero Dirichlet boundary condition. It is clear that the smoothness of $x \mapsto u_\varepsilon^*(t, \omega, x)$ depends on the smoothness of the mappings $x \mapsto g^k(t, \omega, x)$, $k \in \mathbb{N}$, as well as on the properties of the nonlinearities F and G . However, even if F and G vanish, and the spatial smoothness of the g^k is high, the Sobolev smoothness of $x \mapsto u_\varepsilon^*(t, \omega, x)$ can be additionally limited by singularities of the spatial derivatives of u_ε^* at the boundary of \mathcal{O} , due to the zero Dirichlet boundary condition. Such corner singularities are a typical example that the spatial L_p -Sobolev regularity of u_ε^* is exceeded by the regularity in the scale of Besov spaces $B_{\tau, \tau}^\alpha(\mathcal{O})$, $\frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}$. In this paper, we present a result on the spatial Besov regularity of the solution u_ε^* to equation $(*_\varepsilon)$ which has the following structure: If

$$u_\varepsilon^* \in L_p([0, T] \times \Omega, \mathcal{P}_T, \lambda \otimes \mathbb{P}; W_p^s(\mathcal{O}))$$

and if $\varepsilon_f, \varepsilon_g > 0$ are small enough and the functions g^k , $k \in \mathbb{N}$, are sufficiently regular, then

$$u_\varepsilon^* \in L_\tau([0, T] \times \Omega, \mathcal{P}_T, \lambda \otimes \mathbb{P}; B_{\tau, \tau}^\alpha(\mathcal{O}))$$

for certain $\alpha > s$ and $\frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}$. Here \mathcal{P}_T is the predictable σ -algebra w.r.t. the filtration $(\mathcal{F}_t)_{t \in [0, T]}$ and λ denotes Lebesgue measure on $[0, T]$. This result becomes even more important in the light of the following recently proven result, see [27]. Roughly speaking, the following has been shown: Consider the two-dimensional case and assume that the domain $\mathcal{O} \subseteq \mathbb{R}^2$ has a polygonal boundary $\partial\mathcal{O}$ such that \mathcal{O} lies on one side of $\partial\mathcal{O}$. Denote by γ_{\max} the largest interior angle at any corner of the boundary $\partial\mathcal{O}$. Consider equation $(*_\varepsilon)$ with $\varepsilon = 0$ and g being sufficiently smooth (see Section 2 for the concrete setting). Then, the regularity of the solution u_0^* in the scale $L_2(\Omega_T; W_2^s(\mathcal{O}))$, $s > 0$, is limited by $s^* = 1 + \pi/\gamma_{\max}$, which is strictly less than 2 if $\gamma_{\max} > \pi$. However, at the same time $u_0^* \in L_\tau(\Omega_T; B_{\tau, \tau}^\alpha(\mathcal{O}))$, $1/\tau = \alpha/2 + 1/2$, for each $\alpha < 2$. With the explanations from above in mind, this result is important for the theoretical foundation of adaptive numerical methods for the approximation of u_ε^* .

The proofs of our results heavily rely on regularity estimates in weighted Sobolev spaces. For linear SPDEs, estimates of this form have already been derived, see [23], as well as [24]. In this paper, we generalize these results to the semilinear case. By following the lines of [2], these new weighted Sobolev estimates can then be used to derive the desired Besov results.

There exists an extensive literature on the Besov regularity of SPDEs, although most of the results are obtained for linear equations. Recently, *semilinear* evolution equations gained center stage. In general, however, the assumptions on the domain and the scale of parameters considered do not fit into our setting. To mention an example, in [32]

non-autonomous semilinear evolution equations in UMD Banach spaces with M-type 2 have been studied (see also [29] for the autonomous case). As an application, it has been shown that, if the boundary of the domain is smooth enough and certain conditions on the coefficients are fulfilled, the solutions of these equations take values in the Besov space $B_{p,p}^s(\mathcal{O})$ with $p \geq 2$ and some $s \in [0, 2]$. One of the main purposes in these works is to gain better Hölder regularity results. Hence, due to the Sobolev embedding theorem, the main interest is on spaces with summability parameter p as large as possible. However, our focus is different. Firstly, we are explicitly interested in domains with non-smooth boundary. In this situation, one of the main ingredients for the application mentioned above, namely the equality $W_2^2(\mathcal{O}) \cap \mathring{W}_2^1(\mathcal{O}) = D(A)$, where $D(A) := \{u \in \mathring{W}_2^1(\mathcal{O}) : Au \in L_2(\mathcal{O})\}$, $A = \Delta = \sum_{\mu=1}^d \frac{\partial^2}{\partial x_\mu^2}$, does not hold any more (see [19], [20], as well as [22]). Secondly, we are interested in the scale $B_{\tau,\tau}^\alpha(\mathcal{O})$, $1/\tau = \alpha/d + 1/p$, $p \geq 2$ fixed, including spaces which are no Banach spaces but quasi-Banach spaces. The parameter τ decreases if α increases and $B_{\tau,\tau}^\alpha(\mathcal{O})$ fails to be a Banach space for $\tau < 1$. While our methods work in this setting, any direct approach requires (at least!) a fully-fledged theory of stochastic integration in quasi-Banach spaces which is not yet available.

Let us emphasize that our result can be extended to more general semilinear equations of the type

$$\left. \begin{aligned} du &= \left(\sum_{i,j=1}^d a^{ij} u_{x_i x_j} + \sum_{i=1}^d b^i u_{x_i} + cu + \varepsilon_f F(u) + f \right) dt \\ &\quad + \sum_{k=1}^{\infty} \left(\sum_{i=1}^d \sigma^{ik} u_{x_i} + \eta^k u + \varepsilon_g G(u)^k + g^k \right) dw_t^k, \end{aligned} \right\} (\bullet_\varepsilon)$$

$$u(0, \cdot) = u_0.$$

Here the coefficients a^{ij} , b^i , c , σ^{ik} , η^k and the free terms f and g^k are random functions depending on t and x , while F and G are as in equation $(*_\varepsilon)$. This extension is possible because our result in Theorem 8 as well as Corollary 10 can be extended to this wider class. Therefore, additionally assumptions on the coefficients, as presented in [23] and [24], have to be fulfilled. Since this mainly adds notational complications, we will focus on equation $(*_\varepsilon)$ and give a short account of how to treat equations of type (\bullet_ε) in the last section.

The paper is organized as follows: In Section 2 we extend the result from [24] concerning the solvability of linear SPDEs in weighted Sobolev spaces to a class of *semilinear* SPDEs. Furthermore, we will prove a weighted Sobolev estimate for this unique solution. Section 3 contains the Besov regularity result (Theorem 11), its proof and an example for an application of this theorem. General equations of type (\bullet_ε) will be treated in Section 4. In the appendix the reader can find the definition of Besov spaces $B_{p,q}^s(\mathcal{O})$ as well as some details on wavelet characterizations of $B_{p,q}^s(\mathbb{R}^d)$ for particular parameter constellations.

Before we start let us fix some **Conventions and Notations** (see [2]). If not otherwise stated, throughout this paper, $\mathcal{O} \subseteq \mathbb{R}^d$ denotes a *bounded* Lipschitz domain. A domain is

called Lipschitz if each point on the boundary $\partial\mathcal{O}$ has a neighbourhood whose intersection with the boundary—after relabeling and reorienting the coordinate axes if necessary—is the graph of a Lipschitz function.

By $\mathcal{D}'(\mathcal{O})$ we denote the space of Schwartz distributions on \mathcal{O} . If not explicitly stated otherwise, all function spaces or spaces of distributions are meant to be spaces of real-valued functions or distributions. If $f \in \mathcal{D}'(\mathcal{O})$ is a generalized function and $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{N}_0^d$ is a multi-index, we write $D^\alpha f = \frac{\partial^{|\alpha|} f}{\partial x_1^{\alpha_1} \dots \partial x_d^{\alpha_d}}$ for the corresponding derivative w.r.t. $x = (x_1, \dots, x_d) \in \mathcal{O}$, where $|\alpha| = \alpha_1 + \dots + \alpha_d$. For $m \in \mathbb{N}_0$, $D^m f = \{D^\alpha f : |\alpha| = m\}$ is the set of all m -th order derivatives of f which is identified with an $\mathbb{R}^{\binom{d+m-1}{m}}$ -valued distribution. Given $p \in [1, \infty)$ and $m \in \mathbb{N}_0$, $W_p^m(\mathcal{O})$ denotes the classical Sobolev space consisting of all (equivalence classes of) measurable functions $f : \mathcal{O} \rightarrow \mathbb{R}$ such that $\|f\|_{W_p^m(\mathcal{O})} = \|f\|_{L_p(\mathcal{O})} + |f|_{W_p^m(\mathcal{O})} = (\int_{\mathcal{O}} |f(x)|^p dx)^{1/p} + \sum_{|\alpha|=m} (\int_{\mathcal{O}} |D^\alpha f(x)|^p dx)^{1/p}$ is finite. For $p \in (1, \infty)$ and $s \in (m, m+1)$, $m \in \mathbb{N}_0$, we define the fractional order Sobolev space $W_p^s(\mathcal{O})$ to be the Besov space $B_{p,p}^s(\mathcal{O})$ introduced in the appendix. (This scale of fractional order Sobolev spaces can also be obtained by real interpolation of $W_p^n(\mathcal{O})$, $n \in \mathbb{N}_0$. One can show that $W_2^n(\mathcal{O}) = B_{2,2}^n(\mathcal{O})$ for all $n \in \mathbb{N}$ and $W_p^n(\mathcal{O}) \subset B_{p,p}^n(\mathcal{O})$ for all $n \in \mathbb{N}$, $p > 2$, see, e.g. [31, Remark 2.3.3/4 and Theorem 4.6.1.(b)] together with [17].) Given any countable index set \mathcal{J} , the space of p -summable sequences indexed by \mathcal{J} is denoted by $\ell_p = \ell_p(\mathcal{J})$ and $|\cdot|_{\ell_p}$ is the respective norm. Usually we have $\ell_p = \ell_p(\mathbb{N})$ but, for instance we may also use the notation $|D^m f(x)|_{\ell_p}^p = \sum_{|\alpha|=m} |D^\alpha f(x)|^p$ for $f \in W_p^m(\mathcal{O})$.

Given a distribution $f \in \mathcal{D}'(\mathcal{O})$ and a smooth and compactly supported test function $\varphi \in \mathcal{C}_0^\infty(\mathcal{O})$, we write $\langle f, \varphi \rangle$ for the application of f to φ . If H is a Hilbert space, then $\langle \cdot, \cdot \rangle_H$ denotes the inner product in H . $\mathcal{M}_T^{2,c}(H, (\mathcal{F}_t))$ is the space of continuous, square integrable, H -valued martingales with respect to the filtration $(\mathcal{F}_t)_{t \in [0, T]}$. For $[0, T] \times \Omega$ we use the shorthand notation Ω_T and

$$\mathcal{P}_T := \sigma(\{(s, t] \times F_s : 0 \leq s < t \leq T, F_s \in \mathcal{F}_s\} \cup \{\{0\} \times F_0 : F_0 \in \mathcal{F}_0\})$$

is the predictable σ -algebra. $\lambda \otimes \mathbb{P}$ is the product measure of Lebesgue measure λ on $([0, T], \mathcal{B}([0, T]))$, where $\mathcal{B}([0, T])$ denotes the Borel σ -algebra on $[0, T]$, and the probability measure \mathbb{P} on (Ω, \mathcal{F}) . Given any measure space (A, \mathcal{A}, m) , any (quasi-)normed space B with (quasi-)norm $\|\cdot\|_B$ and any summability index $p > 0$, we denote by $L_p(A, \mathcal{A}, m; B)$ the space of all strongly measurable, i.e. Bochner measurable, functions $u : A \rightarrow B$ whose (quasi-)norm $\|u\|_{L_p(A, \mathcal{A}, m; B)} := (\int_A \|u(z)\|_B^p m(dz))^{1/p}$ is finite.

All equalities of random variables or random (generalized) functions appearing in this paper are meant to be \mathbb{P} -almost sure equalities. Throughout the paper, C denotes a positive constant which may change its value from line to line.

2 Existence Results in Weighted Sobolev Spaces

In this chapter we extend the L_p -theory of linear SPDEs on Lipschitz domains from [24] to *semilinear* SPDEs of the form $(*_\varepsilon)$.

If we assume that our solution $u_\varepsilon^* = u_\varepsilon^*(t, \omega, x)$, $(t, \omega, x) \in [0, T] \times \Omega \times \mathcal{O}$, vanishes at the boundary $\partial\mathcal{O}$, satisfying a zero Dirichlet boundary condition, we are facing the same problem as in the linear case: Nonsmooth domains \mathcal{O} induce singularities of the (spatial) derivatives at the boundary $\partial\mathcal{O}$, so that our solution fails to have high regularity in the Sobolev scale $W_p^s(\mathcal{O})$, $s > 0$. A general way to deal with smoothness regardless of certain singularities at the boundary is to use weighted Sobolev spaces, where the weight function is a power of the distance to the boundary. The L_p -theory in [23, 24] is based on spaces of this type, namely the weighted Sobolev spaces $H_{p,\theta}^\gamma(\mathcal{O})$, $p \in (1, \infty)$, $\theta, \gamma \in \mathbb{R}$, introduced in [28]. For the convenience of the reader we give here a short presentation of these spaces, which can also be found in [2]. They are defined in terms of the Bessel-potential spaces

$$H_p^\gamma(\mathbb{R}^d) = \{u \in \mathcal{S}'(\mathbb{R}^d) : \|u\|_{H_p^\gamma(\mathbb{R}^d)} = \|(1 - \Delta)^{\gamma/2} u\|_{L_p(\mathbb{R}^d)} < \infty\}.$$

Here, $\mathcal{S}'(\mathbb{R}^d) \subset \mathcal{D}'(\mathbb{R}^d)$ is the space of (real valued) tempered distributions and $(1 - \Delta)^{\gamma/2} : \mathcal{S}'(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$ is the pseudo-differential operator with symbol $\mathbb{R}^d \ni \xi \mapsto (1 + |\xi|^2)^{\gamma/2}$, i.e. $(1 - \Delta)^{\gamma/2} u = \mathcal{F}^{-1}((1 + |\xi|^2)^{\gamma/2} \mathcal{F}u)$, where \mathcal{F} denotes the Fourier transform on the (complex valued) tempered distributions.

For $x \in \mathcal{O}$ we write $\rho(x) := \text{dist}(x, \partial\mathcal{O})$ for the distance between x and the boundary of the domain \mathcal{O} . Fix $c > 1$, $k_0 > 0$ and for $n \in \mathbb{Z}$ consider the subsets \mathcal{O}_n of \mathcal{O} given by

$$\mathcal{O}_n := \{x \in \mathcal{O} : c^{-n-k_0} < \rho(x) < c^{-n+k_0}\}.$$

Let ζ_n , $n \in \mathbb{Z}$, be non-negative functions satisfying $\zeta_n \in \mathcal{C}_0^\infty(\mathcal{O}_n)$, $\sum_{n \in \mathbb{Z}} \zeta_n(x) = 1$ and $|D^m \zeta_n(x)| \leq C \cdot c^{mn}$ for all $n \in \mathbb{Z}$, $m \in \mathbb{N}_0$, $x \in \mathcal{O}$, and a constant $C > 0$ that does not depend on n , m and x . The functions ζ_n can be constructed by mollifying the indicator functions of the sets \mathcal{O}_n , see, e.g. [21, Section 1.4]. If \mathcal{O}_n is empty we set $\zeta_n \equiv 0$. For $u \in \mathcal{D}'(\mathcal{O})$ $\zeta_n u$ is a distribution on \mathcal{O} with compact support which can be extended by zero to \mathbb{R}^d . This extension is a tempered distribution, i.e. $\zeta_n u \in \mathcal{S}'(\mathbb{R}^d)$.

Definition 1. Let ζ_n , $n \in \mathbb{Z}$, be as above and $p \in (1, \infty)$, $\theta, \gamma \in \mathbb{R}$. Then

$$H_{p,\theta}^\gamma(\mathcal{O}) := \left\{ u \in \mathcal{D}'(\mathcal{O}) : \|u\|_{H_{p,\theta}^\gamma(\mathcal{O})}^p := \sum_{n \in \mathbb{Z}} c^{n\theta} \|\zeta_{-n}(c^n \cdot) u(c^n \cdot)\|_{H_p^\gamma(\mathbb{R}^d)}^p < \infty \right\}.$$

According to [28] this definition is independent of the specific choice of c , k_0 and ζ_n , $n \in \mathbb{N}_0$, in the sense that one gets equivalent norms. If $\gamma = m \in \mathbb{N}_0$ then the spaces can be characterized as

$$\begin{aligned} H_{p,\theta}^0(\mathcal{O}) &= L_{p,\theta}(\mathcal{O}) := L_p(\mathcal{O}, \rho(x)^{\theta-d} dx), \\ H_{p,\theta}^m(\mathcal{O}) &= \{u : \rho^{|\alpha|} D^\alpha u \in L_{p,\theta}(\mathcal{O}) \text{ for all } \alpha \in \mathbb{N}_0^d \text{ with } |\alpha| \leq m\}, \end{aligned}$$

and one has the norm equivalence

$$C^{-1} \|u\|_{H_{p,\theta}^m(\mathcal{O})}^p \leq \sum_{\alpha \in \mathbb{N}_0^d, |\alpha| \leq m} \int_{\mathcal{O}} |\rho(x)^{|\alpha|} D^\alpha u(x)|^p \rho(x)^{\theta-d} dx \leq C \|u\|_{H_{p,\theta}^m(\mathcal{O})}^p.$$

Analogous notations are used for $\ell_2 = \ell_2(\mathbb{N})$ -valued functions $g = (g^k)_{k \in \mathbb{N}}$. For $p \in (1, \infty)$, $\theta, \gamma \in \mathbb{R}$ and $\zeta_n, n \in \mathbb{Z}$, as above

$$\begin{aligned} H_p^\gamma(\mathbb{R}^d; \ell_2) &:= \left\{ g \in (\mathcal{S}'(\mathbb{R}^d))^{\mathbb{N}} : (1 - \Delta)^{\gamma/2} g^k \in L_p(\mathbb{R}^d) \text{ for all } k \in \mathbb{N} \text{ and} \right. \\ &\quad \left. \|g\|_{H_p^\gamma(\mathbb{R}^d; \ell_2)} := \left\| \left\| (1 - \Delta)^{\gamma/2} g^k \right\|_{L_p(\mathbb{R}^d)} \right\|_{\ell_2} < \infty \right\}, \\ H_{p,\theta}^\gamma(\mathcal{O}; \ell_2) &:= \left\{ g \in (\mathcal{D}'(\mathcal{O}))^{\mathbb{N}} : \|g\|_{H_{p,\theta}^\gamma(\mathcal{O}; \ell_2)}^p := \sum_{n \in \mathbb{Z}} c^{n\theta} \|\zeta_{-n}(c^n \cdot) g(c^n \cdot)\|_{H_p^\gamma(\mathbb{R}^d; \ell_2)}^p < \infty \right\}. \end{aligned}$$

Remark 2 (see Remark 7.(a) in [2]). One can consider the spaces $H_{p,\theta}^\gamma(\mathcal{O})$ as generalizations of the classical Sobolev spaces on \mathcal{O} with zero Dirichlet boundary conditions. For $\gamma = m \in \mathbb{N}_0$ we have the identity

$$H_{p,d-mp}^m(\mathcal{O}) = \mathring{W}_p^m(\mathcal{O}),$$

and the norms in both spaces are equivalent, see Theorem 9.7 in [26]. Here $\mathring{W}_p^m(\mathcal{O})$ is the closure of $\mathcal{C}_0^\infty(\mathcal{O})$ in the classical Sobolev space $W_p^m(\mathcal{O})$.

We can now define spaces of stochastic processes and random functions in terms of the weighted Sobolev spaces introduced above.

Definition 3. For $\gamma, \theta \in \mathbb{R}$ and $p \in (1, \infty)$ we set

$$\begin{aligned} \mathbb{H}_{p,\theta}^\gamma(\mathcal{O}, T) &:= L_p(\Omega_T, \mathcal{P}_T, \lambda \otimes \mathbb{P}; H_{p,\theta}^\gamma(\mathcal{O})), \\ \mathbb{H}_{p,\theta}^\gamma(\mathcal{O}, T; \ell_2) &:= L_p(\Omega_T, \mathcal{P}_T, \lambda \otimes \mathbb{P}; H_{p,\theta}^\gamma(\mathcal{O}; \ell_2)), \\ U_{p,\theta}^\gamma(\mathcal{O}) &:= L_p(\Omega, \mathcal{F}_0, \mathbb{P}; H_{p,\theta+2-p}^{\gamma-2/p}(\mathcal{O})), \end{aligned}$$

and for $p \in [2, \infty)$,

$$\begin{aligned} \mathfrak{H}_{p,\theta}^\gamma(\mathcal{O}, T) &:= \left\{ u \in \mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T) : u(0, \cdot) \in U_{p,\theta}^\gamma(\mathcal{O}) \text{ and } du = f dt + \sum_{k=1}^{\infty} g^k dw_t^k \right. \\ &\quad \left. \text{for some } f \in \mathbb{H}_{p,\theta+p}^{\gamma-2}(\mathcal{O}, T), g \in \mathbb{H}_{p,\theta}^{\gamma-1}(\mathcal{O}, T; \ell_2) \right\}, \end{aligned}$$

equipped with the norm

$$\|u\|_{\mathfrak{H}_{p,\theta}^\gamma(\mathcal{O}, T)} := \|u\|_{\mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)} + \|f\|_{\mathbb{H}_{p,\theta+p}^{\gamma-2}(\mathcal{O}, T)} + \|g\|_{\mathbb{H}_{p,\theta}^{\gamma-1}(\mathcal{O}, T; \ell_2)} + \|u(0, \cdot)\|_{U_{p,\theta}^\gamma(\mathcal{O})}.$$

The equality $du = f dt + \sum_{k=1}^{\infty} g^k dw_t^k$ above is a shorthand for

$$\langle u(t, \cdot), \varphi \rangle = \langle u(0, \cdot), \varphi \rangle + \int_0^t \langle f(s, \cdot), \varphi \rangle ds + \sum_{k=1}^{\infty} \int_0^t \langle g^k(s, \cdot), \varphi \rangle dw_s^k \quad (1)$$

for all $\varphi \in \mathcal{C}_0^\infty(\mathcal{O})$, $t \in [0, T]$.

Remark 4 (see Remark 9 in [2]). **(a)** If $p \in [2, \infty)$, then the sum of stochastic integrals in (1) converges in the space $\mathcal{M}_T^{2,c}(\mathbb{R}, (\mathcal{F}_t))$ of continuous, square integrable, \mathbb{R} -valued martingales w.r.t $(\mathcal{F}_t)_{t \in [0, T]}$. A proof can be found in [2, Appendix A].

(b) Using the arguments from [25, Remark 3.3], we get the uniqueness (up to indistinguishability) of the pair $(f, g) \in \mathbb{H}_{p, \theta+p}^{\gamma-2}(\mathcal{O}, T) \times \mathbb{H}_{p, \theta}^{\gamma-1}(\mathcal{O}, T; \ell_2)$ which fulfils (1). Consequently, the norm in $\mathfrak{H}_{p, \theta}^\gamma(\mathcal{O}, T)$ is well defined.

We are now ready to get more concrete on our model equation $(*_\varepsilon)$. In the sequel we will fix

$$p \in [2, \infty), \gamma \in \mathbb{N} \text{ and } \theta \in \mathbb{R}$$

and assume that the following conditions are fulfilled.

Assumptions 5. The differential operator A , the noise term $(g^k)_{k \in \mathbb{N}}$, the nonlinearity F , the initial condition u_0 , and the nonlinearity G in equation $(*_\varepsilon)$ fulfil the following conditions:

[A1] The operator

$$\begin{aligned} A : H_{p, \theta-p}^\gamma(\mathcal{O}) &\rightarrow H_{p, \theta+p}^{\gamma-2}(\mathcal{O}) \\ u &\mapsto Au \end{aligned}$$

is the unique linear and continuous extension of the linear operator

$$\begin{aligned} \Delta : \mathcal{C}_0^\infty(\mathcal{O}) &\rightarrow H_{p, \theta+p}^{\gamma-2}(\mathcal{O}) \\ u &\mapsto \Delta u := \sum_{j=1}^d \frac{\partial^2}{\partial x_j^2} u \end{aligned}$$

on $H_{p, \theta-p}^\gamma(\mathcal{O})$. We will make abuse of notation and denote A by Δ as well.

[A2] The free noise term has the following regularity: $g = (g^k)_{k \in \mathbb{N}} \in \mathbb{H}_{p, \theta}^{\gamma-1}(\mathcal{O}, T; \ell_2)$.

[A3] The function

$$\begin{aligned} F : \mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T) &\rightarrow \mathbb{H}_{p, \theta+p}^{\gamma-2}(\mathcal{O}, T) \\ u &\mapsto F(u) \end{aligned}$$

is Lipschitz continuous with Lipschitz constant $C_F \in (0, \infty)$, i.e.

$$\|F(u) - F(v)\|_{\mathbb{H}_{p, \theta+p}^{\gamma-2}(\mathcal{O}, T)} \leq C_F \|u - v\|_{\mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T)} \quad (2)$$

for arbitrary $u, v \in \mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T)$.

[A4] The function F is zero at the origin, i.e. $F(0) = 0$.

[A5] The initial value has the following regularity: $u_0 \in U_{p,\theta}^\gamma(\mathcal{O})$.

[A6] The function

$$\begin{aligned} G : \mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T) &\rightarrow \mathbb{H}_{p,\theta}^{\gamma-1}(\mathcal{O}, T; \ell_2) \\ u &\mapsto G(u) = (G(u)^k)_{k \in \mathbb{N}} \end{aligned}$$

is Lipschitz continuous with Lipschitz constant $C_G \in (0, \infty)$, i.e.

$$\|G(u) - G(v)\|_{\mathbb{H}_{p,\theta}^{\gamma-1}(\mathcal{O}, T; \ell_2)} \leq C_G \|u - v\|_{\mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)} \quad (3)$$

for arbitrary $u, v \in \mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)$.

[A7] The function G is zero at the origin, i.e. $G(0) = 0$.

Remark 6. (a) To keep notation simple, we restrict ourself in this section to the case $A = \Delta$, see [A1]. In Section 4 we allow more general operators A , including for example positive definite second order elliptic differential operators.

(b) By making slightly abuse of notation we will also denote by Δ the operator

$$\begin{aligned} \tilde{A} : \mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T) &\rightarrow \mathbb{H}_{p,\theta+p}^{\gamma-2}(\mathcal{O}, T) \\ u &\mapsto \tilde{A}u, \end{aligned}$$

where $\tilde{A}u(t, \omega) := \Delta u(t, \omega, \cdot)$ for all $(t, \omega) \in [0, T] \times \Omega$.

Let us now recall what is meant by a *solution* of equation $(*_\varepsilon)$ in the class $\mathfrak{H}_{p,\theta}^\gamma(\mathcal{O}, T)$.

Definition 7. A solution of equation $(*_\varepsilon)$ in the class $\mathfrak{H}_{p,\theta}^\gamma(\mathcal{O}, T)$ is a stochastic process $u_\varepsilon^* \in \mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)$, which fulfils the following equality for all $\varphi \in \mathcal{C}_0^\infty(\mathcal{O})$ and all $t \in [0, T]$ \mathbb{P} -a.s.

$$\begin{aligned} \langle u_\varepsilon^*(t, \cdot), \varphi \rangle &= \langle u_0, \varphi \rangle + \int_0^t \langle \Delta u_\varepsilon^*(s, \cdot) + \varepsilon_f F(u_\varepsilon^*)(s, \cdot), \varphi \rangle ds \\ &\quad + \sum_{k=1}^{\infty} \int_0^t \langle \varepsilon_g G(u_\varepsilon^*)^k(s, \cdot) + g^k(s, \cdot), \varphi \rangle dw_s^k. \end{aligned}$$

We can now prove the following result concerning the existence and uniqueness of a solution to equation $(*_\varepsilon)$.

Theorem 8. Fix $p \in [2, \infty)$ and $\gamma \in \mathbb{N}$. There exists a constant $\kappa = \kappa(d, p, \mathcal{O}) \in (0, 1)$, such that the following holds: If assumptions [A1]–[A7] are satisfied for some $\theta \in (d + p - 2 - \kappa, d + p - 2 + \kappa)$, there exist two constants $a > 0$ and $b > 0$ depending on $d, p, \gamma, \theta, T, \mathcal{O}, C_F$ and C_G , so that for each

$$\varepsilon = (\varepsilon_f, \varepsilon_g) \in B = B(a, b) := \{(x, y) \in [0, \infty) \times [0, \infty) \mid y < a - bx\},$$

equation $(*_\varepsilon)$ has a unique solution u_ε^* in the class $\mathfrak{H}_{p,\theta}^\gamma(\mathcal{O}, T)$. Moreover, the following inequality holds

$$\|u_\varepsilon^*\|_{\mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)} \leq C (\|g\|_{\mathbb{H}_{p,\theta}^{\gamma-1}(\mathcal{O}, T; \ell_2)} + \|u_0\|_{U_{p,\theta}^\gamma(\mathcal{O})}), \quad (4)$$

where the constant $C \in (0, \infty)$ depends only on $d, p, \gamma, \theta, T, \mathcal{O}, \varepsilon, C_F$ and C_G .

Remark 9. (a) The existence of a constant $\kappa = \kappa(d, p, \mathcal{O}) \in (0, 1)$, such that the statements of Theorem 8 are true for the special case of linear equations, i.e. $\varepsilon = (\varepsilon_f, \varepsilon_g) = (0, 0)$, has been proven in [24]. The constant κ in Theorem 8 is the same as the constant β_0 in Theorem 2.12 in [24], see also Remark 2.13 therein, while the constant $C = C(d, p, \gamma, \theta, T, \mathcal{O}, \varepsilon_f, \varepsilon_g, C_F, C_G)$ in (4) is given by

$$C = \frac{C_0}{1 - C_0(C_F\varepsilon_f + C_G\varepsilon_g)},$$

with $C_0 = C(d, p, \gamma, \theta, T, \mathcal{O})$ from [24, Theorem 2.12].

(b) Because Theorem 2.12 in [24] holds true for fractional and negative parameters γ , see Remark 2.13 therein, the statement of Theorem 8 holds true for arbitrary $\gamma \in \mathbb{R}$.

(c) If the nonlinearity in the noise part of equation $(*_\varepsilon)$ vanishes, i.e. $G \equiv 0$, we can choose the Lipschitz constant $C_G \in (0, \infty)$ from assumption [A6] to be as small as we want. Therefore, instead of the set B in Theorem 8, we get the existence of a constant $a > 0$ depending on $d, p, \gamma, \theta, T, \mathcal{O}$ and C_F , such that the rest of the statement is true for each $\varepsilon \in [0, a) \times [0, \infty)$.

Proof of Theorem 8. Fix $p \in [2, \infty)$ and $\gamma \in \mathbb{N}$. Set $\kappa = \beta_0(d, p, \mathcal{O}) \in (0, 1)$ as in [24, Theorem 2.12] and choose $\theta \in (d + p - 2 - \kappa, d + p - 2 + \kappa)$. First we prove the existence and uniqueness of the solution. To achieve this goal, we use an appropriate version of the Banach fixed point theorem. For $\varepsilon = (\varepsilon_f, \varepsilon_g) \in [0, \infty) \times [0, \infty)$, let us define the following operator:

$$\begin{aligned} \mathcal{N}_\varepsilon : \mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T) &\rightarrow \mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T) \\ u &\mapsto \mathcal{N}_\varepsilon(u), \end{aligned}$$

where $\mathcal{N}_\varepsilon(u)$ is the unique solution of the equation

$$dv = (\Delta v + \varepsilon_f F(u)) dt + (\varepsilon_g G(u)^k + g^k) dw_t^k, \quad v(0, \cdot) = u_0$$

in the class $\mathfrak{H}_{p, \theta}^\gamma(\mathcal{O}, T) \subseteq \mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T)$. Note that this operator is well defined due to Theorem 2.12 in [24], see also Remark 2.13 therein. Let us fix two processes $u, v \in \mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T)$. Then, we know that, if we fix $\varphi \in \mathcal{C}_0^\infty(\mathcal{O})$, for all $t \in [0, T]$, \mathbb{P} -a.s.

$$\begin{aligned} \langle \mathcal{N}_\varepsilon(u)(t, \cdot), \varphi \rangle &= \langle u_0, \varphi \rangle + \int_0^t \langle \Delta \mathcal{N}_\varepsilon(u)(s, \cdot) + \varepsilon_f F(u)(s, \cdot), \varphi \rangle ds \\ &\quad + \sum_{k=1}^{\infty} \int_0^t \langle \varepsilon_g G(u)^k(s, \cdot) + g^k(s, \cdot), \varphi \rangle dw_s^k, \end{aligned}$$

as well as

$$\langle \mathcal{N}_\varepsilon(v)(t, \cdot), \varphi \rangle = \langle u_0, \varphi \rangle + \int_0^t \langle \Delta \mathcal{N}_\varepsilon(v)(s, \cdot) + \varepsilon_f F(v)(s, \cdot), \varphi \rangle ds$$

$$+ \sum_{k=1}^{\infty} \int_0^t \langle \varepsilon_g G(v)^k(s, \cdot) + g^k(s, \cdot), \varphi \rangle dw_s^k.$$

Subtracting the two equations gives us

$$\begin{aligned} \langle \mathcal{N}_\varepsilon(u)(t, \cdot) - \mathcal{N}_\varepsilon(v)(t, \cdot), \varphi \rangle &= \int_0^t \langle \Delta \mathcal{N}_\varepsilon(u)(s, \cdot) - \Delta \mathcal{N}_\varepsilon(v)(s, \cdot), \varphi \rangle ds \\ &\quad + \int_0^t \langle \varepsilon_f F(u)(s, \cdot) - \varepsilon_f F(v)(s, \cdot), \varphi \rangle ds \\ &\quad + \sum_{k=1}^{\infty} \int_0^t \langle \varepsilon_g G(u)^k(s, \cdot) - \varepsilon_g G(v)^k(s, \cdot), \varphi \rangle dw_s^k \end{aligned}$$

for all $t \in [0, T]$ \mathbb{P} -a.s. Because the Laplacian is a linear operator, we get

$$\begin{aligned} \langle \mathcal{N}_\varepsilon(u)(t, \cdot) - \mathcal{N}_\varepsilon(v)(t, \cdot), \varphi \rangle &= \int_0^t \langle \Delta(\mathcal{N}_\varepsilon(u)(s, \cdot) - \mathcal{N}_\varepsilon(v)(s, \cdot)), \varphi \rangle ds \\ &\quad + \int_0^t \langle \varepsilon_f F(u)(s, \cdot) - \varepsilon_f F(v)(s, \cdot), \varphi \rangle ds \\ &\quad + \sum_{k=1}^{\infty} \int_0^t \langle \varepsilon_g G(u)^k(s, \cdot) - \varepsilon_g G(v)^k(s, \cdot), \varphi \rangle dw_s^k. \end{aligned}$$

Consequently, this means that $\mathcal{N}_\varepsilon(u) - \mathcal{N}_\varepsilon(v) \in \mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T)$ is the unique solution in the class $\mathfrak{H}_{p, \theta}^\gamma(\mathcal{O}, T)$ of the linear (stochastic) partial differential equation

$$d\tilde{v} = (\Delta \tilde{v} + \varepsilon_f F(u) - \varepsilon_f F(v))dt + (\varepsilon_g G(u)^k - \varepsilon_g G(v)^k)dw_t^k, \quad \tilde{v}(0, \cdot) = 0,$$

and the inequality

$$\begin{aligned} \|\mathcal{N}_\varepsilon(u) - \mathcal{N}_\varepsilon(v)\|_{\mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T)} \\ \leq C_0 \left(\varepsilon_f \|F(u) - F(v)\|_{\mathbb{H}_{p, \theta+p}^{\gamma-2}(\mathcal{O}, T)} + \varepsilon_g \|G(u) - G(v)\|_{\mathbb{H}_{p, \theta}^{\gamma-1}(\mathcal{O}, T; \ell_2)} \right) \end{aligned}$$

holds with some constant $C_0 \in (0, \infty)$ depending only on d, p, γ, θ, T and \mathcal{O} . This follows from Theorem 2.12 in [24], see also Remark 2.13 therein for the case $\gamma = 1$. The main point is that the constant C there does not depend on the free terms f and g , meaning that the constant C_0 in our situation does not depend on F, G, u or v . Now, using assumptions [A3] and [A6], we get

$$\begin{aligned} \|\mathcal{N}_\varepsilon(u) - \mathcal{N}_\varepsilon(v)\|_{\mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T)} &\leq C_0 \left(C_F \varepsilon_f \|u - v\|_{\mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T)} + C_G \varepsilon_g \|u - v\|_{\mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T)} \right) \\ &\leq C_0 (C_F \varepsilon_f + C_G \varepsilon_g) \|u - v\|_{\mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T)}. \end{aligned}$$

It follows that, for

$$\varepsilon = (\varepsilon_f, \varepsilon_g) \in B := \left\{ (x, y) \in [0, \infty) \times [0, \infty) \mid y < \frac{1}{C_0 C_G} - \frac{C_F}{C_G} x \right\},$$

the operator \mathcal{N}_ε is a contraction from $\mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)$ to $\mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)$. Because the space $\mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)$ of stochastic processes is a Banach space, the well-known Banach fixed point theorem gives us the existence of a unique fixed point of \mathcal{N}_ε . Regarding the fact that any solution of the equation $(*_\varepsilon)$ in $\mathfrak{H}_{p,\theta}^\gamma(\mathcal{O}, T)$ is a fixed point of the operator \mathcal{N}_ε and vice versa, we have just proven that for each $\varepsilon \in B \subseteq [0, \infty) \times [0, \infty)$, equation $(*_\varepsilon)$ has a unique solution u_ε^* in the class $\mathfrak{H}_{p,\theta}^\gamma(\mathcal{O}, T)$.

Now, let us prove inequality (4). Therefore, fix $\varepsilon \in B$ and denote by $q_\varepsilon \in (0, 1)$ the contraction number of \mathcal{N}_ε . That is,

$$\|\mathcal{N}_\varepsilon(u) - \mathcal{N}_\varepsilon(v)\|_{\mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)} \leq q_\varepsilon \|u - v\|_{\mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)} \text{ for all } u, v \in \mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T).$$

Let us start our fixed point iteration with $u^{(1)} = 0 \in \mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)$ and set $u^{(j+1)} := \mathcal{N}_\varepsilon(u^{(j)})$ for all $j \in \mathbb{N}$. We know

(*) that the sequence $(u^{(j)})_{j \in \mathbb{N}}$ converges to the solution u_ε^* in $\mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)$, and

(*) that

$$\|\mathcal{N}_\varepsilon(u^{(1)})\|_{\mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O})} \leq C_0 (\|g\|_{\mathbb{H}_{p,\theta}^{\gamma-1}(\mathcal{O}, T; \ell_2)} + \|u_0\|_{U_{p,\theta}^\gamma(\mathcal{O})}).$$

This follows from Theorem 2.12 in [24] and assumptions [A4] and [A7]. See also Remark 2.13 therein for the case $\gamma = 1$.

The a-priori estimate from the Banach fixed point theorem leads to

$$\|u_\varepsilon^*\|_{\mathbb{H}_{p,\theta-p}^\gamma(\mathcal{O}, T)} \leq \frac{1}{1 - q_\varepsilon} C_0 (\|g\|_{\mathbb{H}_{p,\theta}^{\gamma-1}(\mathcal{O}, T; \ell_2)} + \|u_0\|_{U_{p,\theta}^\gamma(\mathcal{O})}).$$

The proof is finished. \square

The following result is a straightforward consequence from Theorem 8. We will need it in the next section where we want to establish Besov regularity of the solution.

Corollary 10. *In the situation of Theorem 8, the following inequality holds for every $\varepsilon \in B$ and $\tau \in [0, p]$.*

$$\int_{\Omega} \int_0^T \|\rho^{\gamma-(1+\frac{d-\theta}{p})} |D^\gamma u_\varepsilon^*(t, \omega, \cdot)|_{\ell_p} \|_{L_p(\mathcal{O})}^\tau dt \mathbb{P}(d\omega) \leq C \left(\|g\|_{\mathbb{H}_{p,\theta}^{\gamma-1}(\mathcal{O}, T; \ell_2)} + \|u_0\|_{U_{p,\theta}^\gamma(\mathcal{O})} \right)^\tau.$$

Proof. Just replace u by u_ε^* in the proof of Corollary 12 in [2] and use Theorem 8 instead of Theorem 11 in [2]. \square

3 Besov Regularity of the Solution

In this section we state and prove the main result of this paper. It is formulated in terms of the L_τ -spaces

$$L_\tau(\Omega_T; B_{\tau,\tau}^s(\mathcal{O})) = L_\tau(\Omega_T, \mathcal{P}_T, \lambda \otimes \mathbb{P}; B_{\tau,\tau}^s(\mathcal{O})), \quad \tau \in (0, \infty), s \in (0, \infty),$$

and the spaces introduced in the last section. We use the convention “ $1/0 := \infty$ ”.

Theorem 11. *Given the situation from Theorem 8, fix $\varepsilon \in B$ and let u_ε^* be the unique solution of equation $(*_\varepsilon)$ in the class $\mathfrak{H}_{p,\theta}^\gamma(\mathcal{O}, T)$. Assume furthermore that*

$$u_\varepsilon^* \in L_p(\Omega_T; B_{p,p}^s(\mathcal{O})) \quad \text{for some } 0 < s \leq \gamma \wedge \left(1 + \frac{d-\theta}{p}\right).$$

Then, we have

$$u_\varepsilon^* \in L_\tau(\Omega_T; B_{\tau,\tau}^\alpha(\mathcal{O})), \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}, \quad \text{for all } 0 < \alpha < \gamma \wedge s \frac{d}{d-1},$$

and the following equation holds

$$\|u_\varepsilon^*\|_{L_\tau(\Omega_T; B_{\tau,\tau}^\alpha(\mathcal{O}))} \leq C \left(\|g\|_{\mathbb{H}_{p,\theta}^{\gamma-1}(\mathcal{O}, T; \ell_2)} + \|u_0\|_{U_{p,\theta}^\gamma(\mathcal{O})} + \|u_\varepsilon^*\|_{L_p(\Omega_T; B_{p,p}^s(\mathcal{O}))} \right).$$

Here the constant $C > 0$ depends only on $d, p, \gamma, \theta, T, \mathcal{O}, \varepsilon, C_F, C_G$ and α .

Remark 12. Since for $p \in (1, \infty)$ and $s \in \mathbb{R} \setminus \mathbb{N}_0$ we have $W_p^s(\mathcal{O}) = B_p^s(\mathcal{O})$, and since $\frac{d}{d-1} > 1$, Theorem 11 implies that the Besov regularity of the solution to $(*_\varepsilon)$ is usually higher than the Sobolev regularity, see also Example 13 below. Consequently, the use of adaptive schemes for the numerical treatment of $(*_\varepsilon)$ is completely justified.

Proof of Theorem 11. We fix $\varepsilon \in B$ and follow the lines of the proof of Theorem 15 in [2]. I.e. we establish Besov regularity by first using extension operators from $B_{p,p}^s(\mathcal{O})$ to $B_{p,p}^s(\mathbb{R}^d)$ as described in [30], and then employing the wavelet characterization according to Theorem 19 in the appendix. The estimation of the wavelet coefficients, which correspond to the wavelets with support inside of the domain \mathcal{O} is performed by the weighted Sobolev results from the previous section. \square

Example 13. Let $p = 2$ and $\gamma = 2$ and choose $\theta = d \in (d-c, d+c)$ for all $c \in (0, 1)$. Furthermore, let $G \equiv 0$ and set

$$\begin{aligned} F : \mathbb{H}_{2,d-2}^2(\mathcal{O}, T) &\rightarrow \mathbb{H}_{2,d+2}^0(\mathcal{O}, T) \\ u &\mapsto F(u), \end{aligned}$$

where for each $u \in \mathbb{H}_{2,d-2}^2(\mathcal{O}, T)$ we set

$$\begin{aligned} F(u) : [0, T] \times \Omega &\rightarrow H_{2,d+2}^0(\mathcal{O}) \\ (t, \omega) &\mapsto F(u)(t, \omega) := \rho^{-2} \sin(u(t, \omega, \cdot)). \end{aligned}$$

Then obviously, $F(0) = 0$ and F is Lipschitz continuous with Lipschitz constant $C_F = 1$. Now, Theorem 8 gives us the existence of an $\varepsilon_{0f} > 0$, such that for any $\varepsilon_f \in (0, \varepsilon_{0f})$ the equation

$$du = (\Delta u + \varepsilon_f F(u))dt + \sum_{k=1}^{\infty} g^k(t)dw_t^k, \quad u(0, \cdot) = u_0$$

has a unique solution u_ε^* in the class $\mathfrak{H}_{2,d}^2(\mathcal{O}, T)$. This implies that our solution also has the following spatial regularity

$$\begin{aligned} u_\varepsilon^* &\in L_2(\Omega_T, \mathcal{P}_T, \lambda \otimes \mathbb{P}; H_{2,d-2}^2(\mathcal{O})) \subseteq L_2(\Omega_T, \mathcal{P}_T, \lambda \otimes \mathbb{P}; H_{2,d-2}^1(\mathcal{O})) \\ &= L_2(\Omega_T, \mathcal{P}_T, \lambda \otimes \mathbb{P}; \mathring{W}_2^1(\mathcal{O})) \\ &\subseteq L_2(\Omega_T, \mathcal{P}_T, \lambda \otimes \mathbb{P}; B_{2,2}^1(\mathcal{O})). \end{aligned}$$

Hence, we can apply Theorem 11 and get

$$u_\varepsilon^* \in L_\tau(\Omega_T, \mathcal{P}_T, \lambda \otimes \mathbb{P}; B_{\tau,\tau}^\alpha(\mathcal{O})) \text{ for all } \alpha \in (0, \frac{d}{d-1}) \text{ with } \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{2}.$$

In the two-dimensional setting ($d = 2$) we conclude for our solution, that

$$u_\varepsilon^* \in L_\tau([0, T] \times \Omega, \mathcal{P}_T, \lambda \otimes \mathbb{P}; B_{\tau,\tau}^\alpha(\mathcal{O})) \text{ for all } \alpha \in (0, 2) \text{ with } \frac{1}{\tau} = \frac{\alpha + 1}{2}.$$

4 A Wider Class of Equations

In the introduction we have indicated that our main result can be extended to equation (\bullet_ε) . The major reason is, that with the same techniques used in the proof of Theorem 8 we can prove existence and uniqueness of a solution to equation (\bullet_ε) , provided the coefficients $a^{ij}, b^i, c, \sigma^{ik}$ and η^k , the free terms f and g^k and the initial value u_0 fulfil certain conditions. This is done by using the general result from [24, Theorem 2.12] and employing again the Banach fixed point theorem. A similar estimate to the one from Corollary 10 follows immediately. We can use this fact to extend our regularity result to such equations. In this section we want to sketch the basic steps.

For the convenience of the reader we begin by presenting the assumptions on the coefficients used in [24, Theorem 2.12]. Therefore, we need some additional notations. For $x, y \in \mathcal{O}$ we shall write $\rho(x, y) := \rho(x) \wedge \rho(y)$. For $\alpha \in \mathbb{R}$, $\delta \in (0, 1]$ and $m \in \mathbb{N}_0$ we set:

$$\begin{aligned} [f]_m^{(\alpha)} &:= \sup_{x \in \mathcal{O}} \rho^{m+\alpha}(x) |D^m f(x)| & \text{and} & & [f]_{m+\delta}^{(\alpha)} &:= \sup_{\substack{x, y \in \mathcal{O} \\ |\beta|=m}} \rho^{m+\alpha}(x, y) \frac{|D^\beta f(x) - D^\beta f(y)|}{|x - y|^\delta}, \\ |f|_m^{(\alpha)} &:= \sum_{j=0}^m [f]_j^{(\alpha)} & \text{and} & & |f|_{m+\delta}^{(\alpha)} &:= |f|_m^{(\alpha)} + [f]_{m+\delta}^{(\alpha)}, \end{aligned}$$

whenever it makes sense. We shall use the same notations for ℓ_2 -valued functions (just replace the absolute values in the above definitions by the ℓ_2 -norms). Furthermore, let's fix an arbitrary function

$$\mu_0 : [0, \infty) \rightarrow [0, \infty),$$

vanishing only on the set of nonnegative integers, i.e. $\mu_0(m) = 0$ if and only if $m \in \mathbb{N}_0$. We set

$$t_+ := t + \mu_0(t).$$

Now we are able to present the assumptions on the coefficients of equation (\bullet_ε) (see [23, Assumptions 2.5 and 2.6] as well as [24, Assumptions 2.10]).

[K1] For any fixed $x \in \mathcal{O}$, the coefficients

$$a^{ij}(\cdot, \cdot, x), b^i(\cdot, \cdot, x), c(\cdot, \cdot, x), \sigma^{ik}(\cdot, \cdot, x), \eta^k(\cdot, \cdot, x) : [0, T] \times \Omega \rightarrow \mathbb{R}$$

are predictable processes with respect to the given normal filtration $(\mathcal{F}_t)_{t \in [0, T]}$.

[K2] (Stochastic parabolicity) There are constants $\delta_0, K > 0$, such that for all $(t, \omega, x) \in [0, T] \times \Omega \times \mathcal{O}$ and $\lambda \in \mathbb{R}^d$:

$$\delta_0 |\lambda|^2 \leq \overline{a^{ij}}(t, \omega, x) \lambda_i \lambda_j \leq K |\lambda|^2,$$

where $\overline{a^{ij}} := a^{ij} - \frac{1}{2} \langle \sigma^i, \sigma^j \rangle_{\ell_2}$ for $i, j \in \{1, \dots, d\}$.

[K3] For all $(t, \omega) \in [0, T] \times \Omega$:

$$\begin{aligned} |a^{ij}(t, \omega, \cdot)|_{|\gamma|_+}^{(0)} + |b^i(t, \omega, \cdot)|_{|\gamma|_+}^{(1)} + |c(t, \omega, \cdot)|_{|\gamma|_+}^{(2)} \\ + |\sigma^i(t, \omega, \cdot)|_{|\gamma|_+}^{(0)} + |\eta(t, \omega, \cdot)|_{|\gamma+1|_+}^{(1)} \leq K. \end{aligned}$$

[K4] The coefficients a^{ij} and σ^i are uniformly continuous in $x \in \mathcal{O}$, i.e. for any $\epsilon > 0$ there is a $\delta = \delta(\epsilon) > 0$, such that

$$|a^{ij}(t, \omega, x) - a^{ij}(t, \omega, y)| + |\sigma^i(t, \omega, x) - \sigma^i(t, \omega, y)|_{\ell_2} \leq \epsilon,$$

for all $(t, \omega) \in [0, T] \times \Omega$, whenever $x, y \in \mathcal{O}$ with $|x - y| \leq \delta$.

[K5] The behaviour of the coefficients b^i , c and η can be controlled near the boundary of \mathcal{O} in the following way:

$$\lim_{\substack{\rho(x) \rightarrow 0 \\ x \in \mathcal{O}}} \sup_{\substack{\omega \in \Omega \\ t \in [0, T]}} \{\rho(x) |b^i(t, \omega, x)| + \rho^2(x) |c(t, \omega, x)| + \rho(x) |\eta(t, \omega, x)|_{\ell_2}\} = 0.$$

Here is the result concerning the solvability of equation (\bullet_ε) in the classes $\mathfrak{H}_{p, \theta}^\gamma(\mathcal{O}, T)$.

Theorem 14. Fix $p \in [2, \infty)$ and $\gamma \in \mathbb{N}$. Furthermore, let assumptions [K1]–[K5] be satisfied for some $K > 0$ and $\delta_0 > 0$. Then there exists a constant $\kappa = \kappa(d, p, \gamma, \delta_0, K, \mathcal{O}) \in (0, 1)$, such that the following holds: If assumptions [A1]–[A7] are satisfied for some $\theta \in (d + p - 2 - \kappa, d + p - 2 + \kappa)$, there exist two constants $a > 0$ and $b > 0$ depending on $d, \gamma, p, \theta, \delta_0, K, T, \mathcal{O}, C_F$ and C_G , so that for each

$$\varepsilon = (\varepsilon_f, \varepsilon_g) \in B = B(a, b) := \{(x, y) \in [0, \infty) \times [0, \infty) \mid y < a - bx\},$$

equation (\bullet_ε) has a unique solution u_ε^\bullet in the class $\mathfrak{H}_{p, \theta}^\gamma(\mathcal{O}, T)$. Moreover, for this solution we have

$$\|u_\varepsilon^\bullet\|_{\mathbb{H}_{p, \theta-p}^\gamma(\mathcal{O}, T)} \leq C \left(\|f\|_{\mathbb{H}_{p, \theta+p}^{\gamma-2}(\mathcal{O}, T)} + \|g\|_{\mathbb{H}_{p, \theta}^{\gamma-1}(\mathcal{O}, T; \ell_2)} + \|u_0\|_{U_{p, \theta}^\gamma(\mathcal{O})} \right), \quad (5)$$

where C is a constant depending only on $d, \gamma, p, \theta, \delta_0, K, T, C_F, C_G, \varepsilon$ and \mathcal{O} .

Remark 15. The statements of Remark 9 (a) and (b) also carry over to Theorem 14.

An immediate consequence of Theorem 14 is the following estimate.

Corollary 16. *In the situation of Theorem 14, the following inequality holds for every $\varepsilon \in B$ and $\tau \in [0, p]$.*

$$\begin{aligned} \int_{\Omega} \int_0^T \|\rho^{\gamma-(1+\frac{d-\theta}{p})} |D^\gamma u_\varepsilon^\bullet(t, \omega, \cdot)|_{\ell_p} \|_{L_p(\mathcal{O})}^\tau dt \mathbb{P}(d\omega) \\ \leq C \left(\|f\|_{\mathbb{H}_{p,\theta+p}^{\gamma-2}(\mathcal{O},T)} + \|g\|_{\mathbb{H}_{p,\theta}^{\gamma-1}(\mathcal{O},T;\ell_2)} + \|u_0\|_{U_{p,\theta}^\gamma(\mathcal{O})} \right)^\tau. \end{aligned}$$

Proof. Just repeat the arguments of the proof of Corollary 10 and use estimate (5) instead of (4) at the beginning. \square

Now we can present our result concerning the Besov regularity of the solution in the generalized setting.

Theorem 17. *Given the situation from Theorem 14, fix $\varepsilon \in B$ and let u_ε^\bullet be the unique solution to equation (\bullet_ε) in the class $\mathfrak{H}_{p,\theta}^\gamma(\mathcal{O},T)$. Furthermore, assume that*

$$u_\varepsilon^\bullet \in L_p(\Omega_T; B_{p,p}^s(\mathcal{O})) \quad \text{for some } 0 < s \leq \gamma \wedge \left(1 + \frac{d-\theta}{p}\right).$$

Then, we have

$$u_\varepsilon^\bullet \in L_\tau(\Omega_T; B_{\tau,\tau}^\alpha(\mathcal{O})), \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}, \quad \text{for all } 0 < \alpha < \gamma \wedge s \frac{d}{d-1},$$

and the following inequality holds

$$\|u_\varepsilon^\bullet\|_{L_\tau(\Omega_T; B_{\tau,\tau}^\alpha(\mathcal{O}))} \leq C \left(\|f\|_{\mathbb{H}_{p,\theta+p}^{\gamma-2}(\mathcal{O},T)} + \|g\|_{\mathbb{H}_{p,\theta}^{\gamma-1}(\mathcal{O},T;\ell_2)} + \|u_0\|_{U_{p,\theta}^\gamma(\mathcal{O})} + \|u_\varepsilon^\bullet\|_{L_p(\Omega_T; B_{p,p}^s(\mathcal{O}))} \right).$$

Proof. We can argue like we did in the proof of Theorem 11. We just have to use Corollary 16 where we used Corollary 10. \square

A Appendix

In this appendix we give a definition of Besov spaces and present a well-known result concerning the wavelet characterization of these function spaces. Our standard reference in this context is the monograph [3].

The most common way to define Besov spaces is by making use of differences. For a function $f : \mathcal{O} \rightarrow \mathbb{R}$ and a natural number $n \in \mathbb{N}$ let

$$\Delta_h^n f(x) := \prod_{i=0}^n \mathbf{1}_{\mathcal{O}}(x + ih) \cdot \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} f(x + jh)$$

be the n -th difference of f with step $h \in \mathbb{R}^d$. For $p \in (0, \infty)$ the modulus of smoothness is given by

$$\omega^n(t, f)_p := \sup_{|h| < t} \|\Delta_h^n f\|_{L_p(\mathcal{O})}, \quad t > 0.$$

One approach to introduce Besov spaces is the following (see, e.g. [16, p.91]).

Definition 18. Let $s, p, q \in (0, \infty)$ and $n \in \mathbb{N}$ with $n > s$. Then $B_{p,q}^s(\mathcal{O})$ is the collection of all functions $f \in L_p(\mathcal{O})$ such that

$$|f|_{B_{p,q}^s(\mathcal{O})} := \left(\int_0^\infty \left[t^{-s} \omega^n(t, f)_p \right]^q \frac{dt}{t} \right)^{1/q} < \infty.$$

These classes are equipped with a (quasi-)norm by taking

$$\|f\|_{B_{p,q}^s(\mathcal{O})} := \|f\|_{L_p(\mathcal{O})} + |f|_{B_{p,q}^s(\mathcal{O})}.$$

For certain constellations of the parameters p, q and s , Besov spaces $B_{p,q}^s(\mathbb{R}^d)$ can be characterized by the decay properties of wavelet coefficients. To this end, fix an arbitrary $r \in \mathbb{N}$, and let $\varphi \in \mathcal{C}^r(\mathbb{R}^d)$ be a compactly supported scaling function as constructed by I. Daubechies, see [15]. We denote by $\psi_i \in \mathcal{C}^r(\mathbb{R}^d)$, $i = 1, \dots, 2^d - 1$, the corresponding multivariate Daubechies mother wavelets having compact support and vanishing moments up to order r . Furthermore, assume that

$$\{\varphi_k, \psi_{i,j,k} : (i, j, k) \in \{1, \dots, 2^d - 1\} \times \mathbb{N} \times \mathbb{Z}^d\}$$

is an orthogonal wavelet basis of $L_2(\mathbb{R}^d)$. (We used the standard abbreviations for dyadic shifts and dilations of functions, see e.g. [2, p.8].) Such a basis can be used for the characterization of Besov spaces as follows.

Theorem 19. Let $p, q \in (0, \infty)$ and $s > \max\{0, d(1/p - 1)\}$. Choose $r \in \mathbb{N}$ with $r > s$ and a corresponding Daubechies orthonormal wavelet basis. Then a locally integrable function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is in the Besov space $B_{p,q}^s(\mathbb{R}^d)$ if, and only if,

$$f = \sum_{k \in \mathbb{Z}^d} \langle f, \varphi_k \rangle \varphi_k + \sum_{i=1}^{2^d-1} \sum_{j \in \mathbb{N}_0} \sum_{k \in \mathbb{Z}^d} \langle f, \psi_{i,j,k} \rangle \psi_{i,j,k}$$

(convergence in $\mathcal{D}'(\mathbb{R}^d)$) with

$$\left(\sum_{k \in \mathbb{Z}^d} |\langle f, \varphi_k \rangle|^p \right)^{1/p} + \left(\sum_{i=1}^{2^d-1} \sum_{j \in \mathbb{N}_0} 2^{jsq} 2^{jd(1/2-1/p)q} \left(\sum_{k \in \mathbb{Z}^d} |\langle f, \psi_{i,j,k} \rangle|^p \right)^{q/p} \right)^{1/q} < \infty, \quad (6)$$

and (6) is an equivalent (quasi-)norm for $B_{p,q}^s(\mathbb{R}^d)$.

References

- [1] P. Binev, W. Dahmen, and R. DeVore, *Adaptive finite element methods with convergence rates*, Numer. Math. 97 (2004), pp. 219–268.
- [2] P.A. Cioica, S. Dahlke, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R.L. Schilling, *Spatial Besov regularity for stochastic partial differential equations on Lipschitz domains*, preprint (2010), submitted for publication. Available at <http://www.dfg-spp1324.de/publications.php>.
- [3] A. Cohen, *Numerical Analysis of Wavelet Methods*, Studies in Mathematics and Its Applications Vol. 32, Elsevier, 2003.
- [4] A. Cohen, W. Dahmen, and R.A. DeVore, *Adaptive wavelet methods for elliptic operator equations: Convergence rates*, Math. Comp. 70 (2001), pp. 27–75.
- [5] A. Cohen, W. Dahmen, and R.A. DeVore, *Adaptive wavelet methods II—Beyond the elliptic case*, Found. Comput. Math. 2 (2002), pp. 203–245.
- [6] A. Cohen, W. Dahmen, and R. DeVore, *Adaptive wavelet methods for nonlinear variational problems*, SIAM J. Numer. Anal. 41 (2003), pp. 1785–1823.
- [7] S. Dahlke, *Besov regularity for elliptic boundary value problems in polygonal domains*, Appl. Math. Lett. 12 (1999), pp. 31–36.
- [8] S. Dahlke, *Besov regularity for interface problems*, ZAMM Z. Angew. Math. Mech. 79 (1999), pp. 383–388.
- [9] S. Dahlke, *Besov regularity for the Neumann problem*, in *Function Spaces, Differential Operators, and Nonlinear Analysis, The Hans Triebel Anniversary Volume*, D. Haroske, T. Runst, and H.-J. Schmeisser, eds., Birkhäuser, Basel-Boston-Berlin, 2003, pp. 267–277.
- [10] S. Dahlke, W. Dahmen, and R.A. DeVore, *Nonlinear approximation and adaptive techniques for solving elliptic operator equations*, in *Multiscale Wavelet Methods for Partial Differential Equations* W. Dahmen, A. Kurdila, and P. Oswald, eds., Academic Press, 1997, pp. 237–283.
- [11] S. Dahlke, W. Dahmen, R. Hochmuth, and R. Schneider, *Stable multiscale bases and local error estimation for elliptic problems*, Appl. Numer. Math. 23 (1997), pp. 21–47.
- [12] S. Dahlke, W. Dahmen, and K. Urban, *Adaptive wavelet methods for saddle point problems—Optimal convergence rates*, SIAM J. Numer. Anal. 40 (2002), pp. 1230–1262.
- [13] S. Dahlke and R.A. DeVore, *Besov regularity for elliptic boundary value problems*, Comm. Partial Differential Equations 22 (1997), pp. 1–16.
- [14] S. Dahlke and W. Sickel, *Besov regularity for the Poisson equation in smooth and polyhedral cones*, in *Sobolev Spaces in Mathematics II, Applications in Analysis and Partial Differential Equations*, V. Maz'ya, ed., International Mathematical Series 9, Springer, jointly published with Tamara Rozhkovskaya Publisher, Novosibirsk, 2008, pp. 123–145.
- [15] I. Daubechies, *Ten Lectures on Wavelets*, SIAM, Philadelphia, 1992.
- [16] R.A. DeVore, *Nonlinear approximation*, Acta Numer. 7 (1998), pp. 51–150.
- [17] S. Dispa, *Intrinsic characterizations of Besov spaces on Lipschitz domains*, Math. Nachr. 260 (2003), pp. 21–33.
- [18] W. Dörfler, *A convergent adaptive algorithm for Poisson's equation*, SIAM J. Numer. Anal. 33 (1996), pp. 1106–1124.
- [19] P. Grisvard, *Elliptic Problems in Nonsmooth Domains*, Monographs and Studies in Mathematics 24, Pitman, Boston-London-Melbourne, 1985.
- [20] P. Grisvard, *Singularities in Boundary Value Problems*, Research Notes in Applied Mathematics 22, Springer, 1992.
- [21] L. Hörmander, *The Analysis of Linear Partial Differential Operators I. Distribution Theory and Fourier Analysis*, 2nd ed., Springer, 1990.
- [22] D. Jerison, C.E. Kenig, *The inhomogeneous Dirichlet problem in Lipschitz domains*, J. Funct. Anal. 130 (1995), pp. 161–219.
- [23] K.-H. Kim, *An L_p -theory of SPDEs on Lipschitz domains*, Potential Anal. 29 (2008), pp. 303–326.

- [24] K.-H. Kim, *A weighted Sobolev space theory of parabolic stochastic PDEs on non-smooth domains*, preprint (2011), submitted for publication. Available at <http://arxiv.org/abs/1109.4727>.
- [25] N.V. Krylov, *An analytic approach to SPDEs*, in *Stochastic Partial Differential Equations. Six Perspectives*, B.L. Rozovskii, R. Carmona, eds., Mathematical Surveys and Monographs 64, American Mathematical Society, 1999, pp. 185–242.
- [26] A. Kufner, *Weighted Sobolev Spaces*, Teubner-Texte zur Mathematik 31, Teubner, Leipzig, 1980.
- [27] F. Lindner, *Approximation and regularity of stochastic partial differential equations*, Dr. diss., TU Dresden, 2011.
- [28] S.V. Lototsky, *Sobolev spaces with weights in domains and boundary value problems for degenerate elliptic equations*, *Methods Appl. Anal.* 7 (2000), pp. 195–204.
- [29] J.M.A.M. van Neerven, M.C. Veraar, and L. Weis, *Stochastic evolution equations in UMD Banach spaces*, *J. Funct. Anal.* 255 (2008), pp. 940–993.
- [30] V.S. Rychkov, *On restrictions and extensions of the Besov and Triebel-Lizorkin spaces with respect to Lipschitz domains*, *J. Lond. Math. Soc.* 60 (1999), pp. 237–257.
- [31] H. Triebel, *Interpolation Theory, Function Spaces, Differential Operators*, 2nd ed., Barth, Leipzig, 1995.
- [32] M.C. Veraar, *Non-autonomous stochastic evolution equations and applications to stochastic partial differential equations*, *J. Evol. Equ.* 10 (2010), pp. 85–127.

Petru A. Cioica, Stephan Dahlke
 Philipps-Universität Marburg
 FB Mathematik und Informatik, AG Numerik/Optimierung
 Hans-Meerwein-Straße
 35032 Marburg
 Germany
 {cioica, dahlke}@mathematik.uni-marburg.de

Preprint Series DFG-SPP 1324

<http://www.dfg-spp1324.de>

Reports

- [1] R. Ramlau, G. Teschke, and M. Zhariy. A Compressive Landweber Iteration for Solving Ill-Posed Inverse Problems. Preprint 1, DFG-SPP 1324, September 2008.
- [2] G. Plonka. The Easy Path Wavelet Transform: A New Adaptive Wavelet Transform for Sparse Representation of Two-dimensional Data. Preprint 2, DFG-SPP 1324, September 2008.
- [3] E. Novak and H. Woźniakowski. Optimal Order of Convergence and (In-) Tractability of Multivariate Approximation of Smooth Functions. Preprint 3, DFG-SPP 1324, October 2008.
- [4] M. Espig, L. Grasedyck, and W. Hackbusch. Black Box Low Tensor Rank Approximation Using Fibre-Crosses. Preprint 4, DFG-SPP 1324, October 2008.
- [5] T. Bonesky, S. Dahlke, P. Maass, and T. Raasch. Adaptive Wavelet Methods and Sparsity Reconstruction for Inverse Heat Conduction Problems. Preprint 5, DFG-SPP 1324, January 2009.
- [6] E. Novak and H. Woźniakowski. Approximation of Infinitely Differentiable Multivariate Functions Is Intractable. Preprint 6, DFG-SPP 1324, January 2009.
- [7] J. Ma and G. Plonka. A Review of Curvelets and Recent Applications. Preprint 7, DFG-SPP 1324, February 2009.
- [8] L. Denis, D. A. Lorenz, and D. Trede. Greedy Solution of Ill-Posed Problems: Error Bounds and Exact Inversion. Preprint 8, DFG-SPP 1324, April 2009.
- [9] U. Friedrich. A Two Parameter Generalization of Lions' Nonoverlapping Domain Decomposition Method for Linear Elliptic PDEs. Preprint 9, DFG-SPP 1324, April 2009.
- [10] K. Bredies and D. A. Lorenz. Minimization of Non-smooth, Non-convex Functionals by Iterative Thresholding. Preprint 10, DFG-SPP 1324, April 2009.
- [11] K. Bredies and D. A. Lorenz. Regularization with Non-convex Separable Constraints. Preprint 11, DFG-SPP 1324, April 2009.

- [12] M. Döhler, S. Kunis, and D. Potts. Nonequispaced Hyperbolic Cross Fast Fourier Transform. Preprint 12, DFG-SPP 1324, April 2009.
- [13] C. Bender. Dual Pricing of Multi-Exercise Options under Volume Constraints. Preprint 13, DFG-SPP 1324, April 2009.
- [14] T. Müller-Gronbach and K. Ritter. Variable Subspace Sampling and Multi-level Algorithms. Preprint 14, DFG-SPP 1324, May 2009.
- [15] G. Plonka, S. Tenorth, and A. Iske. Optimally Sparse Image Representation by the Easy Path Wavelet Transform. Preprint 15, DFG-SPP 1324, May 2009.
- [16] S. Dahlke, E. Novak, and W. Sickel. Optimal Approximation of Elliptic Problems by Linear and Nonlinear Mappings IV: Errors in L_2 and Other Norms. Preprint 16, DFG-SPP 1324, June 2009.
- [17] B. Jin, T. Khan, P. Maass, and M. Pidcock. Function Spaces and Optimal Currents in Impedance Tomography. Preprint 17, DFG-SPP 1324, June 2009.
- [18] G. Plonka and J. Ma. Curvelet-Wavelet Regularized Split Bregman Iteration for Compressed Sensing. Preprint 18, DFG-SPP 1324, June 2009.
- [19] G. Teschke and C. Borries. Accelerated Projected Steepest Descent Method for Nonlinear Inverse Problems with Sparsity Constraints. Preprint 19, DFG-SPP 1324, July 2009.
- [20] L. Grasedyck. Hierarchical Singular Value Decomposition of Tensors. Preprint 20, DFG-SPP 1324, July 2009.
- [21] D. Rudolf. Error Bounds for Computing the Expectation by Markov Chain Monte Carlo. Preprint 21, DFG-SPP 1324, July 2009.
- [22] M. Hansen and W. Sickel. Best m-term Approximation and Lizorkin-Triebel Spaces. Preprint 22, DFG-SPP 1324, August 2009.
- [23] F.J. Hickernell, T. Müller-Gronbach, B. Niu, and K. Ritter. Multi-level Monte Carlo Algorithms for Infinite-dimensional Integration on \mathbb{R}^N . Preprint 23, DFG-SPP 1324, August 2009.
- [24] S. Dereich and F. Heidenreich. A Multilevel Monte Carlo Algorithm for Lévy Driven Stochastic Differential Equations. Preprint 24, DFG-SPP 1324, August 2009.
- [25] S. Dahlke, M. Fornasier, and T. Raasch. Multilevel Preconditioning for Adaptive Sparse Optimization. Preprint 25, DFG-SPP 1324, August 2009.

- [26] S. Dereich. Multilevel Monte Carlo Algorithms for Lévy-driven SDEs with Gaussian Correction. Preprint 26, DFG-SPP 1324, August 2009.
- [27] G. Plonka, S. Tenorth, and D. Roşca. A New Hybrid Method for Image Approximation using the Easy Path Wavelet Transform. Preprint 27, DFG-SPP 1324, October 2009.
- [28] O. Koch and C. Lubich. Dynamical Low-rank Approximation of Tensors. Preprint 28, DFG-SPP 1324, November 2009.
- [29] E. Faou, V. Gradinaru, and C. Lubich. Computing Semi-classical Quantum Dynamics with Hagedorn Wavepackets. Preprint 29, DFG-SPP 1324, November 2009.
- [30] D. Conte and C. Lubich. An Error Analysis of the Multi-configuration Time-dependent Hartree Method of Quantum Dynamics. Preprint 30, DFG-SPP 1324, November 2009.
- [31] C. E. Powell and E. Ullmann. Preconditioning Stochastic Galerkin Saddle Point Problems. Preprint 31, DFG-SPP 1324, November 2009.
- [32] O. G. Ernst and E. Ullmann. Stochastic Galerkin Matrices. Preprint 32, DFG-SPP 1324, November 2009.
- [33] F. Lindner and R. L. Schilling. Weak Order for the Discretization of the Stochastic Heat Equation Driven by Impulsive Noise. Preprint 33, DFG-SPP 1324, November 2009.
- [34] L. Kämmerer and S. Kunis. On the Stability of the Hyperbolic Cross Discrete Fourier Transform. Preprint 34, DFG-SPP 1324, December 2009.
- [35] P. Cerejeiras, M. Ferreira, U. Kähler, and G. Teschke. Inversion of the noisy Radon transform on $SO(3)$ by Gabor frames and sparse recovery principles. Preprint 35, DFG-SPP 1324, January 2010.
- [36] T. Jahnke and T. Udrescu. Solving Chemical Master Equations by Adaptive Wavelet Compression. Preprint 36, DFG-SPP 1324, January 2010.
- [37] P. Kittipoom, G. Kutyniok, and W.-Q Lim. Irregular Shearlet Frames: Geometry and Approximation Properties. Preprint 37, DFG-SPP 1324, February 2010.
- [38] G. Kutyniok and W.-Q Lim. Compactly Supported Shearlets are Optimally Sparse. Preprint 38, DFG-SPP 1324, February 2010.
- [39] M. Hansen and W. Sickel. Best m -Term Approximation and Tensor Products of Sobolev and Besov Spaces – the Case of Non-compact Embeddings. Preprint 39, DFG-SPP 1324, March 2010.

- [40] B. Niu, F.J. Hickernell, T. Müller-Gronbach, and K. Ritter. Deterministic Multi-level Algorithms for Infinite-dimensional Integration on \mathbb{R}^N . Preprint 40, DFG-SPP 1324, March 2010.
- [41] P. Kittipoom, G. Kutyniok, and W.-Q Lim. Construction of Compactly Supported Shearlet Frames. Preprint 41, DFG-SPP 1324, March 2010.
- [42] C. Bender and J. Steiner. Error Criteria for Numerical Solutions of Backward SDEs. Preprint 42, DFG-SPP 1324, April 2010.
- [43] L. Grasedyck. Polynomial Approximation in Hierarchical Tucker Format by Vector-Tensorization. Preprint 43, DFG-SPP 1324, April 2010.
- [44] M. Hansen und W. Sickel. Best m -Term Approximation and Sobolev-Besov Spaces of Dominating Mixed Smoothness - the Case of Compact Embeddings. Preprint 44, DFG-SPP 1324, April 2010.
- [45] P. Binev, W. Dahmen, and P. Lamby. Fast High-Dimensional Approximation with Sparse Occupancy Trees. Preprint 45, DFG-SPP 1324, May 2010.
- [46] J. Ballani and L. Grasedyck. A Projection Method to Solve Linear Systems in Tensor Format. Preprint 46, DFG-SPP 1324, May 2010.
- [47] P. Binev, A. Cohen, W. Dahmen, R. DeVore, G. Petrova, and P. Wojtaszczyk. Convergence Rates for Greedy Algorithms in Reduced Basis Methods. Preprint 47, DFG-SPP 1324, May 2010.
- [48] S. Kestler and K. Urban. Adaptive Wavelet Methods on Unbounded Domains. Preprint 48, DFG-SPP 1324, June 2010.
- [49] H. Yserentant. The Mixed Regularity of Electronic Wave Functions Multiplied by Explicit Correlation Factors. Preprint 49, DFG-SPP 1324, June 2010.
- [50] H. Yserentant. On the Complexity of the Electronic Schrödinger Equation. Preprint 50, DFG-SPP 1324, June 2010.
- [51] M. Guillemard and A. Iske. Curvature Analysis of Frequency Modulated Manifolds in Dimensionality Reduction. Preprint 51, DFG-SPP 1324, June 2010.
- [52] E. Herrholz and G. Teschke. Compressive Sensing Principles and Iterative Sparse Recovery for Inverse and Ill-Posed Problems. Preprint 52, DFG-SPP 1324, July 2010.
- [53] L. Kämmerer, S. Kunis, and D. Potts. Interpolation Lattices for Hyperbolic Cross Trigonometric Polynomials. Preprint 53, DFG-SPP 1324, July 2010.

- [54] G. Kutyniok and W.-Q Lim. Shearlets on Bounded Domains. Preprint 54, DFG-SPP 1324, July 2010.
- [55] A. Zeiser. Wavelet Approximation in Weighted Sobolev Spaces of Mixed Order with Applications to the Electronic Schrödinger Equation. Preprint 55, DFG-SPP 1324, July 2010.
- [56] G. Kutyniok, J. Lemvig, and W.-Q Lim. Compactly Supported Shearlets. Preprint 56, DFG-SPP 1324, July 2010.
- [57] A. Zeiser. On the Optimality of the Inexact Inverse Iteration Coupled with Adaptive Finite Element Methods. Preprint 57, DFG-SPP 1324, July 2010.
- [58] S. Jokar. Sparse Recovery and Kronecker Products. Preprint 58, DFG-SPP 1324, August 2010.
- [59] T. Aboiyar, E. H. Georgoulis, and A. Iske. Adaptive ADER Methods Using Kernel-Based Polyharmonic Spline WENO Reconstruction. Preprint 59, DFG-SPP 1324, August 2010.
- [60] O. G. Ernst, A. Mugler, H.-J. Starkloff, and E. Ullmann. On the Convergence of Generalized Polynomial Chaos Expansions. Preprint 60, DFG-SPP 1324, August 2010.
- [61] S. Holtz, T. Rohwedder, and R. Schneider. On Manifolds of Tensors of Fixed TT-Rank. Preprint 61, DFG-SPP 1324, September 2010.
- [62] J. Ballani, L. Grasedyck, and M. Kluge. Black Box Approximation of Tensors in Hierarchical Tucker Format. Preprint 62, DFG-SPP 1324, October 2010.
- [63] M. Hansen. On Tensor Products of Quasi-Banach Spaces. Preprint 63, DFG-SPP 1324, October 2010.
- [64] S. Dahlke, G. Steidl, and G. Teschke. Shearlet Coorbit Spaces: Compactly Supported Analyzing Shearlets, Traces and Embeddings. Preprint 64, DFG-SPP 1324, October 2010.
- [65] W. Hackbusch. Tensorisation of Vectors and their Efficient Convolution. Preprint 65, DFG-SPP 1324, November 2010.
- [66] P. A. Cioica, S. Dahlke, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Spatial Besov Regularity for Stochastic Partial Differential Equations on Lipschitz Domains. Preprint 66, DFG-SPP 1324, November 2010.

- [67] E. Novak and H. Woźniakowski. On the Power of Function Values for the Approximation Problem in Various Settings. Preprint 67, DFG-SPP 1324, November 2010.
- [68] A. Hinrichs, E. Novak, and H. Woźniakowski. The Curse of Dimensionality for Monotone and Convex Functions of Many Variables. Preprint 68, DFG-SPP 1324, November 2010.
- [69] G. Kutyniok and W.-Q Lim. Image Separation Using Shearlets. Preprint 69, DFG-SPP 1324, November 2010.
- [70] B. Jin and P. Maass. An Analysis of Electrical Impedance Tomography with Applications to Tikhonov Regularization. Preprint 70, DFG-SPP 1324, December 2010.
- [71] S. Holtz, T. Rohwedder, and R. Schneider. The Alternating Linear Scheme for Tensor Optimisation in the TT Format. Preprint 71, DFG-SPP 1324, December 2010.
- [72] T. Müller-Gronbach and K. Ritter. A Local Refinement Strategy for Constructive Quantization of Scalar SDEs. Preprint 72, DFG-SPP 1324, December 2010.
- [73] T. Rohwedder and R. Schneider. An Analysis for the DIIS Acceleration Method used in Quantum Chemistry Calculations. Preprint 73, DFG-SPP 1324, December 2010.
- [74] C. Bender and J. Steiner. Least-Squares Monte Carlo for Backward SDEs. Preprint 74, DFG-SPP 1324, December 2010.
- [75] C. Bender. Primal and Dual Pricing of Multiple Exercise Options in Continuous Time. Preprint 75, DFG-SPP 1324, December 2010.
- [76] H. Harbrecht, M. Peters, and R. Schneider. On the Low-rank Approximation by the Pivoted Cholesky Decomposition. Preprint 76, DFG-SPP 1324, December 2010.
- [77] P. A. Cioica, S. Dahlke, N. Döhring, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Adaptive Wavelet Methods for Elliptic Stochastic Partial Differential Equations. Preprint 77, DFG-SPP 1324, January 2011.
- [78] G. Plonka, S. Tenorth, and A. Iske. Optimal Representation of Piecewise Hölder Smooth Bivariate Functions by the Easy Path Wavelet Transform. Preprint 78, DFG-SPP 1324, January 2011.
- [79] A. Mugler and H.-J. Starkloff. On Elliptic Partial Differential Equations with Random Coefficients. Preprint 79, DFG-SPP 1324, January 2011.

- [80] T. Müller-Gronbach, K. Ritter, and L. Yaroslavtseva. A Derandomization of the Euler Scheme for Scalar Stochastic Differential Equations. Preprint 80, DFG-SPP 1324, January 2011.
- [81] W. Dahmen, C. Huang, C. Schwab, and G. Welper. Adaptive Petrov-Galerkin methods for first order transport equations. Preprint 81, DFG-SPP 1324, January 2011.
- [82] K. Grella and C. Schwab. Sparse Tensor Spherical Harmonics Approximation in Radiative Transfer. Preprint 82, DFG-SPP 1324, January 2011.
- [83] D.A. Lorenz, S. Schiffler, and D. Tiede. Beyond Convergence Rates: Exact Inversion With Tikhonov Regularization With Sparsity Constraints. Preprint 83, DFG-SPP 1324, January 2011.
- [84] S. Dereich, M. Scheutzow, and R. Schottstedt. Constructive quantization: Approximation by empirical measures. Preprint 84, DFG-SPP 1324, January 2011.
- [85] S. Dahlke and W. Sickel. On Besov Regularity of Solutions to Nonlinear Elliptic Partial Differential Equations. Preprint 85, DFG-SPP 1324, January 2011.
- [86] S. Dahlke, U. Friedrich, P. Maass, T. Raasch, and R.A. Ressel. An adaptive wavelet method for parameter identification problems in parabolic partial differential equations. Preprint 86, DFG-SPP 1324, January 2011.
- [87] A. Cohen, W. Dahmen, and G. Welper. Adaptivity and Variational Stabilization for Convection-Diffusion Equations. Preprint 87, DFG-SPP 1324, January 2011.
- [88] T. Jahnke. On Reduced Models for the Chemical Master Equation. Preprint 88, DFG-SPP 1324, January 2011.
- [89] P. Binev, W. Dahmen, R. DeVore, P. Lamby, D. Savu, and R. Sharpley. Compressed Sensing and Electron Microscopy. Preprint 89, DFG-SPP 1324, March 2011.
- [90] P. Binev, F. Blanco-Silva, D. Blom, W. Dahmen, P. Lamby, R. Sharpley, and T. Vogt. High Quality Image Formation by Nonlocal Means Applied to High-Angle Annular Dark Field Scanning Transmission Electron Microscopy (HAADF-STEM). Preprint 90, DFG-SPP 1324, March 2011.
- [91] R. A. Ressel. A Parameter Identification Problem for a Nonlinear Parabolic Differential Equation. Preprint 91, DFG-SPP 1324, May 2011.
- [92] G. Kutyniok. Data Separation by Sparse Representations. Preprint 92, DFG-SPP 1324, May 2011.

- [93] M. A. Davenport, M. F. Duarte, Y. C. Eldar, and G. Kutyniok. Introduction to Compressed Sensing. Preprint 93, DFG-SPP 1324, May 2011.
- [94] H.-C. Kreuzler and H. Yserentant. The Mixed Regularity of Electronic Wave Functions in Fractional Order and Weighted Sobolev Spaces. Preprint 94, DFG-SPP 1324, June 2011.
- [95] E. Ullmann, H. C. Elman, and O. G. Ernst. Efficient Iterative Solvers for Stochastic Galerkin Discretizations of Log-Transformed Random Diffusion Problems. Preprint 95, DFG-SPP 1324, June 2011.
- [96] S. Kunis and I. Melzer. On the Butterfly Sparse Fourier Transform. Preprint 96, DFG-SPP 1324, June 2011.
- [97] T. Rohwedder. The Continuous Coupled Cluster Formulation for the Electronic Schrödinger Equation. Preprint 97, DFG-SPP 1324, June 2011.
- [98] T. Rohwedder and R. Schneider. Error Estimates for the Coupled Cluster Method. Preprint 98, DFG-SPP 1324, June 2011.
- [99] P. A. Cioica and S. Dahlke. Spatial Besov Regularity for Semilinear Stochastic Partial Differential Equations on Bounded Lipschitz Domains. Preprint 99, DFG-SPP 1324, July 2011.