


DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

Error bound for piecewise deterministic processes modeling stochastic reaction systems

T. Jahnke, M. Kreim

Preprint 119



Edited by

AG Numerik/Optimierung
Fachbereich 12 - Mathematik und Informatik
Philipps-Universität Marburg
Hans-Meerwein-Str.
35032 Marburg


DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

Error bound for piecewise deterministic processes modeling stochastic reaction systems

T. Jahnke, M. Kreim

Preprint 119



The consecutive numbering of the publications is determined by their chronological order.

The aim of this preprint series is to make new research rapidly available for scientific discussion. Therefore, the responsibility for the contents is solely due to the authors. The publications will be distributed by the authors.

Error bound for piecewise deterministic processes modeling stochastic reaction systems*

Tobias Jahnke[†] and Michael Kreim[†]

March 29, 2012

Abstract

Biological processes involving the random interaction of d species with integer particle numbers are often modeled by a Markov jump process on \mathbb{N}_0^d . Realizations of this process can, in principle, be generated with the classical stochastic simulation algorithm proposed in [19], but for very reactive systems this method is usually inefficient. Hybrid models based on piecewise deterministic processes offer an attractive alternative which can decrease the simulation time considerably in applications where species with rather low particle numbers interact with very abundant species. We investigate the convergence of the hybrid model to the original one for a class of reaction systems with two distinct scales. Our main result is an error bound which states that, under suitable assumptions, the hybrid model approximates the marginal distribution of the discrete species and the conditional moments of the continuous species up to an error of $\mathcal{O}(M^{-1})$ where M is the scaling parameter of the partial thermodynamic limit.

Keywords: Stochastic reaction systems, Gillespie algorithm, hybrid models, piecewise deterministic processes, error bounds, chemical master equation, thermodynamic limit

AMS subject classifications: 60J22, 60J27, 65C20, 65C40, 92-08, 92C42, 92D25

1 Introduction

Many processes inside living cells are characterized by low particle numbers and a high degree of randomness which brings about stochastic effects such as, e.g., the switching of a genetic toggle switch or random extinction [34, 33, 13, 39]. Such effects are crucial for the understanding of the biological system but can usually not be reproduced with the traditional deterministic reaction-rate approach where the concentrations of the species are modeled by a set of ordinary differential equations. Often much better results are obtained when the system is considered as a continuous-time Markov jump process on the state space

*Supported by the “Concept for the Future” of Karlsruhe Institute of Technology within the framework of the German Excellence Initiative, and the DFG priority programme SPP 1324 “Mathematical methods for extracting quantifiable information from complex systems”.

[†]Karlsruhe Institute of Technology, Department of Mathematics, Institute for Applied and Numerical Mathematics, Kaiserstr. 93, D-76133 Karlsruhe, Germany, tobias.jahnke@kit.edu, michael.kreim@kit.edu.

of vectors of particle numbers, because this approach respects both the discreteness of the populations and the randomness of the dynamics; cf. [19, 18, 26, 16, 3]. Unbiased realizations of the Markov process can be generated with the stochastic simulation algorithm derived in D. Gillespie’s seminal paper [19]. Although this algorithm is widely used, its efficiency is constrained by the fact that each single reaction event requires an update of the system, which makes the simulation of highly reactive systems very costly.

This has motivated the construction of many new numerical methods which either reformulate the original scheme in a more economical way (cf. [17]) or reduce the simulation time at the price of a small approximation error; cf. [22, 5, 36, 32, 11, 12, 29, 2] and references therein. A second group of approaches aims to simplify the *model* by assuming, e.g., that a part of the reaction system is nearly in equilibrium [35, 7] or that the highly reactive (“fast”) part of the problem can be described in terms of ordinary or stochastic differential equations which can be solved more efficiently; cf. [1, 6, 24, 23, 4, 38]. Such *hybrid models* reduce the numerical costs significantly in many applications, but a rigorous analytical investigation of the error caused by simplifying the model is missing in most cases.

The aim of this article is to prove an error bound for one of these hybrid models. In the investigated approach, a Markov jump process describing the dynamics of the species with low particle numbers is coupled to ordinary differential equations representing the highly abundant species, which removes the fast time-scales from the system and makes numerical simulations (e.g. by a Strang splitting method) much faster. Such an approach is called a *piecewise deterministic model* in the literature (cf. [10, 9, 8, 4, 37, 40]) because between two jumps of the Markov jump process the other part of the systems simply evolves according to an ordinary differential equation. Intuitively it can be expected that in a *partial thermodynamic limit* (i.e. when the abundance of the “large” species tends to infinity but the reaction constants are rescaled in a way to be explained below) the hybrid system converges to the original Markov jump process. This conjecture has recently been confirmed in [8] where convergence of the process in distribution was proved. Although the objective of our work is very similar, our analysis follows a rather different approach. In the error bound presented in Theorem 1 we only consider the error in the marginal distribution of the species described by the Markov process and the conditional expectations of the other species. In exchange, we prove not only convergence but also determine the *rate* of convergence. In fact, it will be shown that the error in the marginal distribution and in the conditional expectations is proportional to M^{-1} where M is the scaling parameter of the partial thermodynamic limit.

This goal is motivated and formulated in more detail in Section 2. As a starting point, stochastic reaction systems modeled by continuous-time Markov jump processes are considered (Section 2.1). The convergence to the classical reaction-rate equations in the thermodynamical limit (Section 2.2) motivates the use of a hybrid model based on a piecewise deterministic process (Section 2.3). Instead of discussing these concepts in an abstract setting, the derivation and the advantage of the piecewise deterministic process is first illustrated by a model problem which serves as an example throughout the paper. In Section 3 we pass from the model problem to a general class of scaled and partitioned reaction systems and define the associated Markov jump process and the corresponding piecewise deterministic process. The probability density of these processes evolve according to the chemical master equation (Section 3.3) and the Liouville-master equation (Section 3.4), and these differential equation will play a central role in the proof of the main result, which is presented

and proved in Section 4.

Throughout the article, most of the constants which appear in inequalities and which are of minor interest are denoted by the same symbol C although the value of these constants may differ from case to case.

2 Motivation

2.1 Modeling reaction systems with Markov processes

It is widely acknowledged that modeling reaction systems by ordinary differential equations (ODEs) yields qualitatively wrong results if the discreteness of the species and/or the stochastic nature of the dynamics cannot be neglected. In these cases, a significantly better approach is to consider the system as a Markov jump process on a discrete state space. This well-known model has been discussed, e.g., in [19, 18, 26, 16, 3], but in order to introduce our notation and to keep the exposition self-contained, we briefly summarize the basic facts.

Consider a discrete number of particles which move randomly in a bounded domain. It is assumed that the conditions of the environment (i.e. the temperature, the size of the domain, etc.) do not change, and that the system is well-stirred in the sense that at any time the random position of each of the particles is uniformly distributed. Each particle belongs to exactly one of $d \in \mathbb{N}$ different species $\mathcal{S}_1, \dots, \mathcal{S}_d$, and particles of the same species are indistinguishable. It is assumed that the position of a single particle does not matter and that only the number of particles of \mathcal{S}_k at time $t \geq 0$, denoted by $X_k(t) \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$, is relevant. The vector $X(t) = (X_1(t), \dots, X_d(t))^T$ of particle numbers changes according to r reaction channels R_1, \dots, R_r which represent events such as, e.g., a reactive collision between two particles from different species which produces a particle of another type, the decomposition of a particle into two particles, the conversion of a particle from one species to another one, the inflow of new particles into the systems from an exterior source, the “death” of particles and so on. Each reaction channel is defined by the equation



with constants $\kappa_{jk}^{in}, \kappa_{jk}^{out} \in \mathbb{N}_0$ and $j \in \{1, \dots, r\}$. If R_j fires at time t , then the particle numbers jump from the old state $X(t^-)$ to the new state $X(t) = X(t^-) + \nu_j$ where

$$\nu_j = \left(\kappa_{j1}^{out} - \kappa_{j1}^{in}, \dots, \kappa_{jd}^{out} - \kappa_{jd}^{in} \right)^T \in \mathbb{Z}^d \quad (2)$$

is called the stoichiometric vector of R_j . The time of the next reaction event and the number of the reaction channel which fires are both random, but depend only on the current state, such that $X(t)$ is a realization of a Markov jump process. To be more precise, we assume that

$$X(t) = n_0 + \sum_{j=1}^r \mathcal{P}_j \left(\int_0^t \alpha_j(X(s)) ds \right) \nu_j \quad (3)$$

with independent Poisson processes \mathcal{P}_j (cf. [16, 4, 2]) and initial state $X(0) = n_0 \in \mathbb{N}^d$. The function α_j is called the propensity of R_j and is typically defined as

$$\alpha_j(n) = c_j \prod_{k=1}^d \binom{n_k}{\kappa_j^k} \quad (4)$$

with a reaction constant $c_j > 0$ which depend on physical properties such as, e.g., the temperature or the volume of the domain. Here and below

$$\binom{x}{k} = \begin{cases} \frac{1}{k!} \prod_{j=0}^{k-1} (x-j) & \text{if } x > k-1 \text{ and } k \in \mathbb{N} \\ 0 & \text{if } x \leq k-1 \text{ and } k \in \mathbb{N} \\ 1 & \text{if } k = 0 \end{cases} \quad (5)$$

denotes the generalized binomial coefficient. For later use we remark that (5) is not restricted to positive integers and can also be evaluated for $x \in \mathbb{R}$.

The probability distribution $p(t, n) = \mathbb{P}(X(t) = n \mid X(0) = n_0)$ associated to $X(t)$ is the solution of the chemical master equation (CME)

$$\begin{aligned} \partial_t p(t, n) &= \sum_{k=1}^r \left(\alpha_k(n - \nu_k) p(t, n - \nu_k) - \alpha_k(n) p(t, n) \right) \\ p(0, n) &= \delta_{n_0}(n) = \begin{cases} 1 & \text{if } n = n_0 \\ 0 & \text{otherwise} \end{cases} \end{aligned} \quad (6)$$

(cf. [20]) with the convention that $p(t, n - \nu_k) = 0$ for all $n - \nu_k \notin \mathbb{N}_0^d$. Solving the CME numerically is a highly non-trivial problem because in most cases (6) has to be solved on a large state space, which excludes all standard methods. Constructing efficient *deterministic* methods for the CME is a new and very active field of research, which, in spite of its importance, is beyond the scope of this work. In this article, the CME will be used for analytical purposes only.

As a typical interpretation of the above model one may consider a *chemical* reaction system where the particles of the species correspond to the molecules of the chemical substances, but the same setting can be used to model, e.g., discrete stochastic predator-prey systems, viral kinetics, and gene regulatory networks.

The stochastic evolution of $X(t)$ can be simulated with the *Stochastic Simulation Algorithm* (SSA) derived in [19]:

1. Let $t = 0$ and $X = n_0$ where $n_0 \in \mathbb{N}_0^d$ is the initial state.
2. Compute $|\alpha(X)| = \sum_{j=1}^r \alpha_j(X)$.
3. Draw two random numbers ϱ_1 and ϱ_2 from the uniform distribution on $[0, 1]$.
4. Compute the time increment: $\tau = \ln(1/\varrho_1)/|\alpha(X)|$,

5. Determine the number of the reaction channel which fires next: find j such that

$$\sum_{k=1}^{j-1} \alpha_k(X) < \varrho_2 |\alpha(X)| \leq \sum_{k=1}^j \alpha_k(X).$$

6. Update the time $t \leftarrow t + \tau$ and the state $X \leftarrow X + \nu_j$.

7. If $t < t_{end}$ go to step 2.

The SSA is very popular among computational biologists but its efficiency is severely affected if the system is highly reactive, i.e. if some of the propensities are large. This happens in particular if the particle numbers of some of the species are relatively high. In this case, the expected value of the time increment τ (i.e. the waiting time between two reaction events) is so small that a huge number of steps is necessary to simulate the system over the full time-interval, and since each step requires two random numbers and an update of the state variable X , the total runtime increases considerably.

2.2 Thermodynamic limit

Before this approach is discussed, we briefly recall that the stochastic model (Markov jump process) converges to the deterministic one (reaction-rate equations, ODEs) in the thermodynamic limit. Consider the *scaled* kinetics

$$X^{[M]}(t) = Mn_0 + \sum_{j=1}^r \mathcal{P}_j \left(\int_0^t \alpha_j^{[M]}(X^{[M]}(s)) ds \right) \nu_j, \quad (7)$$

$$\alpha_j^{[M]}(n) = M^{(1-|\kappa_j^{in}|_1)} \alpha_j(n), \quad |\kappa_j^{in}|_1 = \sum_{i=1}^d \kappa_{ji}^{in} \quad (8)$$

where $M \in \mathbb{N}$ is a scaling parameter which can be thought of as the volume of the domain in which the particles move. If M increases, then the initial particle numbers $X^{[M]}(0) = Mn_0$ and thus the reactivity of the system increase, too. At the same time, the propensities (8) are rescaled in such a way that for $n \in \mathbb{N}_0^d$ one obtains

$$\begin{aligned} \alpha_j^{[M]}(Mn) &= c_j M^{(1-|\kappa_j^{in}|_1)} \prod_{k=1}^d \binom{Mn_k}{\kappa_{jk}^{in}} \\ &\approx c_j M^{(1-|\kappa_j^{in}|_1)} \prod_{k=1}^d \frac{(Mn_k)^{\kappa_{jk}^{in}}}{\kappa_{jk}^{in}!} = c_j M \prod_{k=1}^d \frac{n_k^{\kappa_{jk}^{in}}}{\kappa_{jk}^{in}!} = \mathcal{O}(M) \end{aligned}$$

for every reaction channel $j = \{1, \dots, r\}$. For $M = 1$, the original process (3) is recovered. The thermodynamic limit is the limit dynamics of the *normalized* process $X^{[M]}(t)/M$ for $M \rightarrow \infty$. In his seminal work [30, 31, 16] T. Kurtz proved that

$$\lim_{M \rightarrow \infty} \mathbb{P} \left(\sup_{s \in [0, t_{end}]} \left| \frac{X^{[M]}(t)}{M} - y(t) \right| > \varepsilon \right) = 0 \quad (9)$$

j	Reaction channel R_j	κ_{j1}^{in}	κ_{j1}^{out}	ν_j	Propensity $\alpha_j(n)$	Propensity $\alpha_j^{[M]}(n)$
1	$R_1 : \star \longrightarrow S$	0	1	1	c_1	Mc_1
2	$R_2 : S + S \longrightarrow S$	2	1	-1	$c_2 \frac{n(n-1)}{2}$	$c_2 \frac{n(n-1)}{2M}$

Table 1: Example I: A reaction system with one species $\mathcal{S} = \mathcal{S}_1$ and two reaction channels R_1 and R_2 . For simplicity, we write n instead of n_1 .

where $y(t)$ is the solution of the classical reaction-rate equation

$$\dot{y}(t) = \sum_{j=1}^r \nu_j c_j \prod_{k=1}^d \frac{y_k^{\kappa_{jk}^{in}}(t)}{\kappa_{jk}^{in}!}, \quad y(0) = n_0. \quad (10)$$

This convergence means, roughly speaking, that for sufficiently large particle numbers the discrete, stochastic model (7) can be replaced by the deterministic ordinary differential equation (10). The advantage is that solving (10) is considerably less expensive than generating a sufficient number of realizations of the highly reactive Markov process (7).

This can be illustrated by the following simple example. Consider one single species $\mathcal{S} = \mathcal{S}_1$ which evolves according to the two reaction channels R_1 and R_2 defined in Table 1. The first reaction channel R_1 represents an inflow of new particles from an infinite source. The growth of the population is limited by R_2 which represents competition among the particles. In this example (7) takes the special form

$$X^{[M]}(t) = Mn_0 + \mathcal{P}_1(tMc_1) - \mathcal{P}_2 \left(c_2 \int_0^t \frac{X^{[M]}(s)(X^{[M]}(s) - 1)}{2M} ds \right), \quad (11)$$

and the normalized process $X^{[M]}(t)/M$ converges to the solution of the reaction-rate equation

$$\frac{dy}{dt}(t) = c_1 - c_2 \frac{y^2(t)}{2}, \quad y(0) = n_0. \quad (12)$$

Equation (12) can be derived from (11) in a loose but intuitively plausible way by dividing (11) by M , by approximating $(X^{[M]}(t) - 1)/M \approx X^{[M]}(t)/M \approx y(t)$, and by approximating

$$\frac{1}{M} \mathcal{P}_1(tMc_1) \approx tc_1 \quad \text{and} \quad \frac{1}{M} \mathcal{P}_2 \left(M \frac{c_2}{2} \int_0^t y^2(s) ds \right) \approx \frac{c_2}{2} \int_0^t y^2(s) ds$$

according to the law of large numbers. This yields

$$y(t) = n_0 + tc_1 - \frac{c_2}{2} \int_0^t y^2(s) ds \quad (13)$$

which is the integral form of the reaction-rate equation (12).

j	Reaction channel R_j	Propensity $\alpha_j^{[M]}(n_1, n_2, n_3)$	Stoich. vector ν_j
1	$R_1 : \mathcal{S}_1 \longrightarrow \mathcal{S}_1 + \mathcal{S}_3$	Mc_1n_1	$(0, 0, 1)^T$
2	$R_2 : \mathcal{S}_1 + \mathcal{S}_3 \longrightarrow \mathcal{S}_2$	$\frac{c_2}{M}n_1n_3$	$(-1, 1, -1)^T$
3	$R_3 : \mathcal{S}_2 \longrightarrow \mathcal{S}_1 + \mathcal{S}_3$	c_3n_2	$(1, -1, 1)^T$
4	$R_4 : \mathcal{S}_3 + \mathcal{S}_3 \longrightarrow \emptyset$	$c_4 \frac{n_3(n_3 - 1)}{2M}$	$(0, 0, -2)^T$

Table 2: Example II: A reaction system with three species and four reactions.

2.3 Partial thermodynamic limit and piecewise deterministic processes

In many biological systems the average particle numbers of the species can differ by several orders of magnitude, and often the dynamics of the abundant species is crucially affected by the stochastic behavior of the species with rather low particle numbers. We thus consider now a more complicated model problem consisting of three species $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3$ which interact via the four reaction channels defined in Table 2. \mathcal{S}_1 represents a gene which catalyzes the production of a protein \mathcal{S}_3 via R_1 . For simplicity, the processes of transcription and translation are merged. The protein can auto-regulate the production of new copies by binding to the promoter region of the gene and blocking the transcription. \mathcal{S}_2 denotes the group of blocked genes, and the binding process is described by R_2 . With a certain probability, however, the protein can be released from the gene via the third reaction channel R_3 . Finally, R_4 describes dimerization of the protein to some irrelevant waste product. Since the waste product is not of any interest and does not have any feedback on $\mathcal{S}_1, \mathcal{S}_2$ and \mathcal{S}_3 , its particle numbers do not have to be computed. This is the reason why this species is called \emptyset instead of \mathcal{S}_4 . Due to the particular structure of the system, the total number of genes $X_1^{[M]}(t) + X_2^{[M]}(t) = X_1^{[M]}(0) + X_2^{[M]}(0) = 10$ is constant. As a consequence, one could easily reduce the system to a two-dimensional problem by expressing $X_2^{[M]}(t)$ in terms of $X_1^{[M]}(t)$ or vice versa, but this does not matter here.

We assume that at $t = 0$ there are 10 open genes and no blocked genes, and that there are much more proteins than open or blocked genes. Thus, the scaling from the previous example is now applied to \mathcal{S}_3 only. To be more precise, we let

$$X^{[M]}(0) = (10, 0, M)^T$$

and scale the propensities as indicated in Table 2. For a general reaction system, the scaling rules will be defined in Section 3. At the moment, we only remark that due to this scaling we have

$$X_1(t) + X_2(t) = 10 \ll \mathbb{E} \left(X_3^{[M]}(t) \right) = \mathcal{O}(M)$$

on bounded time intervals, which means that the expected particle numbers lie on two clearly separated scales. For this reaction system, (3) takes the form

$$\begin{aligned}
X^{[M]}(t) = & \begin{pmatrix} 10 \\ 0 \\ M \end{pmatrix} + \mathcal{P}_1 \left(\int_0^t M c_1 X_1^{[M]}(s) ds \right) \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \\
& + \mathcal{P}_2 \left(\int_0^t \frac{c_2}{M} X_1^{[M]}(s) X_3^{[M]}(s) ds \right) \begin{pmatrix} -1 \\ 1 \\ -1 \end{pmatrix} \\
& + \mathcal{P}_3 \left(\int_0^t c_3 X_2^{[M]}(s) ds \right) \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \\
& + \mathcal{P}_4 \left(\int_0^t c_4 \frac{X_3^{[M]}(s)(X_3^{[M]}(s) - 1)}{2M} ds \right) \begin{pmatrix} 0 \\ 0 \\ -2 \end{pmatrix}.
\end{aligned} \tag{14}$$

Now the example from 2.2 suggests that the abundant species \mathcal{S}_3 could be represented by a traditional reaction-rate equation while \mathcal{S}_1 and \mathcal{S}_2 should still be described in the discrete and stochastic setting. Approximating

$$y(s) \approx \frac{X_3^{[M]}(s)}{M} \approx \frac{X_3^{[M]}(s) - 1}{M}$$

and proceeding as in the previous example leads to

$$\begin{aligned}
Z^{[M]}(t) = & \begin{pmatrix} 10 \\ 0 \end{pmatrix} + \mathcal{P}_2 \left(\int_0^t c_2 Z_1(s) y^{[M]}(s) ds \right) \begin{pmatrix} -1 \\ 1 \end{pmatrix} \\
& + \mathcal{P}_3 \left(\int_0^t c_3 Z_2(s) ds \right) \begin{pmatrix} 1 \\ -1 \end{pmatrix}
\end{aligned} \tag{15}$$

$$\begin{aligned}
y^{[M]}(t) = & 1 + \int_0^t c_1 Z_1^{[M]}(s) ds - \frac{1}{M} \int_0^t c_2 Z_1^{[M]}(s) y^{[M]}(s) ds \\
& + \frac{1}{M} \int_0^t c_3 Z_2^{[M]}(s) ds - \int_0^t c_4 \left(y^{[M]}(s) \right)^2 ds
\end{aligned} \tag{16}$$

where $Z_i^{[M]}(s) \approx X_i^{[M]}(s)$ for $i \in \{1, 2\}$. In contrast to the previous example, the equation for $y^{[M]}(t)$ is *not* a purely deterministic ODE since the stochastic variables Z_1 and Z_2 appear on the right-hand side of (16). However, these variables only jump at discrete times and remain constant in between. If no jump occurs in the time interval $[t_1, t_2]$, then there are constants $z_1, z_2 \in \mathbb{N}_0$ such that $Z_1(s) = z_1$ and $Z_2(s) = z_2$ for all $s \in [t_1, t_2]$, and (16) is thus equivalent to the ODE

$$\frac{dy^{[M]}(t)}{dt} = c_1 z_1 - \frac{c_2 z_1}{M} y^{[M]}(t) + \frac{c_3 z_2}{M} - c_4 \left(y^{[M]}(t) \right)^2 \tag{17}$$

for $t \in [t_1, t_2]$. This explains why such approaches are called *piecewise deterministic models* in the literature, see, e.g., [10, 9, 8, 37, 40] and references therein.

The piecewise deterministic model (15)-(16) can be simulated, e.g., with a stochastic Strang splitting method:

1. Choose $t_{\text{end}} > 0$ and $N \in \mathbb{N}$. Let $h = t_{\text{end}}/N$ and $t_i = ih$ for all $i = 0, \dots, N$.
2. Initialize: Set $i = 0$, $Z_0 = (10, 0)$, $y_0 = M$.
3. Half-step in Z :
Perform a stochastic simulation of (15) on the interval $[t_i, t_i + h/2]$ with initial state Z_i and while keeping $y^{[M]}(s) \equiv y_i$ constant. This can be done with SSA or approximate methods such as, e.g., tau-leaping [22]. Let $Z_{i+\frac{1}{2}}$ be the result of this step.
4. Full step in y :
Solve the ordinary differential equation (17) on the interval $[t_i, t_{i+1}]$ with initial value y_i while keeping $(z_1, z_2) = Z_{i+\frac{1}{2}}$ fixed. This can be done, e.g., with a Runge-Kutta or multi-step method. Let y_{i+1} be the result of this step.
5. Half-step in Z :
Perform a stochastic simulation of (15) on the interval $[t_i + h/2, t_{i+1}]$ with initial state $Z_{i+\frac{1}{2}}$ and while keeping $y^{[M]}(s) \equiv y_{i+1}$ constant at the updated value. Let Z_{i+1} be the result of this step.
6. If $i < N$: set $i \leftarrow i + 1$ and go to step 3.

The values Z_i and y_i are approximations to $Z^{[M]}(t_i)$ and $y(t_i)$, respectively. In addition to the *model error* which arises when the exact process $X^{[M]}(t)$ is replaced by the hybrid approach (15)-(16), there is a *splitting error* due to the fact that the two parts of the system are not propagated simultaneously, but in an alternating way. It is well-known, however, that under suitable regularity conditions the Strang splitting converges in distribution with order 2 to the exact solution; see, e.g. [29]. The main topic of this article is an analysis of the *model error*, and we will henceforth assume that the time-step h is so small that the *numerical error* can be neglected.

In order to investigate the accuracy of the piecewise deterministic process (15)-(16), the splitting method was applied to the model problem with $M = 5, 10, 20, 30, 40, 50, 100$ and

$$c_1 = 0.5, c_2 = 3, c_3 = 1, c_4 = 5, t_{\text{end}} = 0.5, N = 100, h = 0.005. \quad (18)$$

For each value of M , 10^6 realizations have been computed, with the exception of $M = 100$ where $2.5 \cdot 10^6$ realizations were generated to increase the accuracy. From this data, the marginal distribution $n_1 \mapsto \mathbb{P}(Z_1^{[M]}(t_{\text{end}}) = n_1)$ was estimated by producing a histogram and dividing by the number of samples. This approximate marginal distribution is not only affected by the model error and the splitting error, but also by the sampling error – the error due to estimating the marginal distribution from a finite number of realizations. But because of the large number of samples and the small step size h , it can be assumed¹ that the total error is dominated by the model error. The approximate marginal distribution was

¹We have checked this by rerunning the experiment with different time-steps and a larger number of samples.

compared with a reference solution which was obtained by solving the CME (6) with high accuracy² and taking sums over n_2 and n_3 . Figure 1 shows a considerable model error for $M = 10$ (left panel). For $M = 50$, however, the piecewise deterministic process yields a very accurate approximation to the true marginal distribution (right panel). The convergence for increasing values of M is illustrated in the left panel of Figure 2. This plot suggests that the error in the marginal distribution is proportional to M^{-1} , and the goal of this article is to prove that this is indeed the case; cf. Theorem 1.

We remark that the model problem considered here is obviously very small. This is unavoidable because a highly accurate reference solution is not available for larger problems. Moreover, in order to investigate the model error, the sampling error has to be reduced by a huge number of realizations, and the required number increases when larger problems are considered. However, our intention is not to demonstrate the efficiency of piecewise deterministic processes nor their applicability to large, complex biological systems; such applications can be found, e.g., in [9, 40]. Our example is only supposed to serve as an illustration of the somewhat abstract analysis in the second part of the paper.




Figure 1: Marginal distributions of the piecewise deterministic model (15)-(16) (thin line) and of the exact process (14) (bold line) for $M = 10$ (left) and $M = 50$ (right) with the configuration given in (18). The marginal distribution $n_1 \mapsto \mathbb{P}(Z_1^{[M]}(t_{\text{end}}) = n_1)$ of the piecewise deterministic model was estimated from the histogram of 1,000,000 samples. The marginal distribution of the exact Markov process was obtained by solving the CME (6) with high accuracy and taking sums over n_2 and n_3 .

²In this special case, the CME can be solved with a standard ODE integrator because the state space is small. As we have pointed out before, such a straightforward computation is not possible when larger reaction systems are investigated.





Figure 2: Illustration of Theorem 1 by the model problem defined in Table 2 with the parameters from (18). The left panel shows the convergence of the marginal distribution of the piecewise deterministic model (15)-(16) to the marginal distribution of the full process (14). The former was estimated from stochastic simulations whereas the latter was obtained by solving the CME (6) with high accuracy and taking sums over n_2 and n_3 . The right panel shows the left-hand side of the inequality (43). In both panels, the comparison with the function $M \mapsto 1/M$ (dotted line) suggests that each of the two error terms decays proportional to $1/M$, which will be confirmed by the error bounds (42) and (43) from Theorem 1.

3 The general setting: two-scale reaction systems

3.1 Reaction channels, propensities and stoichiometry

Motivated by the previous section we consider reaction systems where the species are divided into two groups $\mathcal{S}_1, \dots, \mathcal{S}_d$ and $\mathcal{S}_{d+1}, \dots, \mathcal{S}_D$ with $d, D \in \mathbb{N}$. The particle numbers of $\mathcal{S}_1, \dots, \mathcal{S}_d$ and $\mathcal{S}_{d+1}, \dots, \mathcal{S}_D$ are denoted by the entries of $n \in \mathbb{N}_0^d$ and $m \in \mathbb{N}_0^D$, respectively. Later, $\mathcal{S}_{d+1}, \dots, \mathcal{S}_D$ are supposed to be represented by real, continuous variables whereas the $\mathcal{S}_1, \dots, \mathcal{S}_d$ will still be described by integer particle numbers. Let the j -th reaction channel be given by



j	Reaction channel R_j	$\alpha_j(n)$	$\beta_j^{[M]}(m)$	$\tilde{\beta}_j^{[M]}(y)$	ν_j	μ_j
1	$R_1 : \mathcal{S}_1 \longrightarrow \mathcal{S}_1 + \mathcal{S}_3$	$c_1 n_1$	M	M	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	1
2	$R_2 : \mathcal{S}_1 + \mathcal{S}_3 \longrightarrow \mathcal{S}_2$	$c_2 n_1$	m/M	y	$\begin{pmatrix} -1 \\ 1 \end{pmatrix}$	-1
3	$R_3 : \mathcal{S}_2 \longrightarrow \mathcal{S}_1 + \mathcal{S}_3$	$c_3 n_2$	1	1	$\begin{pmatrix} 1 \\ -1 \end{pmatrix}$	1
4	$R_4 : \mathcal{S}_3 + \mathcal{S}_3 \longrightarrow \emptyset$	c_4	$\frac{m(m-1)}{2M}$	$M \frac{y^2}{2}$	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	-2

Table 3: Example II revisited. In this example, we have $d = 2$, $D = 1$, $r = 4$, $J_0 = \{2, 3\}$, $J_1 = \{1, 4\}$.

with $\kappa_{jk}^{in}, \kappa_{jk}^{out}, \lambda_{jk}^{in}, \lambda_{jk}^{out} \in \mathbb{N}_0$. The stoichiometric vectors are $(\nu_j, \mu_j) \in \mathbb{Z}^{d+D}$ with

$$\begin{aligned} \nu_j &= \left(\kappa_{j1}^{out} - \kappa_{j1}^{in}, \dots, \kappa_{jd}^{out} - \kappa_{jd}^{in} \right)^T \in \mathbb{Z}^d \\ \mu_j &= \left(\lambda_{j1}^{out} - \lambda_{j1}^{in}, \dots, \lambda_{jd}^{out} - \lambda_{jd}^{in} \right)^T \in \mathbb{Z}^D. \end{aligned} \quad (20)$$

For every $j = 1, \dots, r$ the propensity of R_j is supposed to have the form $\alpha_j(n)\beta_j^{(M)}(m)$ where the entries of $n \in \mathbb{N}_0^d$ and $m \in \mathbb{N}_0^D$ are the particle numbers of $\mathcal{S}_1, \dots, \mathcal{S}_d$ and $\mathcal{S}_{d+1}, \dots, \mathcal{S}_D$, respectively. The function $\alpha_j(n)$ is defined by (4) as before. In order to define $\beta_j^{(M)}(m)$, we distinguish two different groups of reactions with index sets

$$J_0 = \{j \in \{1, \dots, r\} : \nu_j = (0, \dots, 0)^T\}, \quad J_1 = \{1, \dots, r\} \setminus J_0; \quad (21)$$

cf. [8]. If $j \in J_0$, then the particle numbers of $\mathcal{S}_1, \dots, \mathcal{S}_d$ are not changed by the j -th reaction channel R_j . Note that this does *not* mean that R_j is independent of these species, since any of the $\mathcal{S}_1, \dots, \mathcal{S}_d$ could act as a catalyst of that reaction; cf. [8]. This is the case, e.g., in the first reaction channel of Table 3. With the indicator function

$$\gamma(j) = \begin{cases} 0 & \text{if } j \in J_0 \\ 1 & \text{if } j \in J_1 \end{cases}. \quad (22)$$

we define

$$\beta_j^{[M]}(x) = M^{1-\gamma(j)} M^{-|\lambda_j^{in}|_1} \prod_{k=1}^D \binom{x_k}{\lambda_{jk}^{in}} \quad (23)$$

for any $x \in \mathbb{R}_+^D$, where $|\lambda_j^{in}|_1 = \sum_{i=1}^d \lambda_{ji}^{in}$.

Example. For the reaction system defined in Table 2 we let $d = 2$ and $D = 1$. Thus, we have $r = 4$, $J_0 = \{1, 4\}$, and $J_1 = \{2, 3\}$. The propensities and the stoichiometric vectors are listed in Table 3; as before, we write m instead of m_1 for convenience.

In most applications, reactions between more than two particles are rather improbable events, because the probability that more than two particles collide is usually negligible.

Therefore, in most reaction systems, we have $|\lambda_j^{in}|_1 + |\kappa_j^{in}|_1 \leq 2$. In the results below, however, a weaker assumption will be made:

Assumption 1 For every $j \in \{1, \dots, r\}$ we assume that $|\lambda_j^{in}|_1 \leq 2$.

The above definitions guarantee that the particle numbers are nonnegative for all times, because a jump to a state outside of $\mathbb{N}_0^d \times \mathbb{N}_0^D$ can never occur, as the following lemma shows.

Lemma 1 For all $j = 1, \dots, r$, the functions α_j and $\beta_j^{[M]}$ have the following property:

1. If $n \in \mathbb{N}_0^d$ but one or several entries of $n + \nu_j$ are negative, then $\alpha_j(n) = 0$.
2. If $m \in \mathbb{N}_0^D$ but one or several entries of $m + \mu_j$ are negative, then $\beta_j^{[M]}(m) = 0$ for every $M \in \mathbb{N}$.

Proof. Suppose that the k -th entry of $n + \nu_j$ is negative. By (20), this means that $n_k + \kappa_{jk}^{out} - \kappa_{jk}^{in} < 0$. Since $n_k, \kappa_{jk}^{in} \in \mathbb{N}_0$ it follows that $n_k \leq \kappa_{jk}^{in} - 1$ and, by definition (4), that $\alpha_j(n) = 0$. The second assertion follows analogously from the definition (23). ■

3.2 Spaces and norms

Henceforth, $\|\cdot\|$ denotes an arbitrary vector norm on \mathbb{R}^N or the induced matrix norm, respectively. For $K \in \mathbb{N}$ let

$$\ell_K^1 = \left\{ u : \mathbb{N}_0^K \longrightarrow \mathbb{R}^N \mid \sum_{n \in \mathbb{N}_0^K} \|u(n)\| < \infty \right\}, \quad \|u\|_{\ell_K^1} = \sum_{n \in \mathbb{N}_0^K} \|u(n)\|$$

be the multivariate and vector-valued counterpart of the standard ℓ^1 -space, which is recovered for $K = N = 1$. In the analysis below only $N = 1$ or $N = d$ will appear, and since the latter case is a generalization of the former, the symbol ℓ_K^1 will be used in both cases. For $i \in \mathbb{N}_0$ we define the spaces $\mathcal{X}_{d,D}^{i+1}$ via the recursion

$$\mathcal{X}_{d,D}^{i+1} = \left\{ u \in \mathcal{X}_{d,D}^i \mid (n, m) \mapsto m_k u(n, m) \in \mathcal{X}_{d,D}^i \text{ for all } k \in \{1, \dots, D\} \right\}$$

with $\mathcal{X}_{d,D}^0 = \ell_{d+D}^1$. If, for example, u is a probability distribution and $u = u(n, m) \in \mathcal{X}_{d,D}^2$, then the first and second moments of the marginal distribution $m \mapsto \sum_{n \in \mathbb{N}_0^d} u(n, m)$ exist.

For later use, we also define the corresponding spaces

$$\begin{aligned} \mathcal{Y}_{d,D}^0 &= \left\{ u : \mathbb{N}_0^d \times \mathbb{R}_+^D \longrightarrow \mathbb{R}^N \mid \sum_{n \in \mathbb{N}_0^d} \int_{\mathbb{R}_+^D} \|u(n, x)\| dx < \infty \right\} \\ \mathcal{Y}_{d,D}^{i+1} &= \left\{ u \in \mathcal{Y}_{d,D}^i \mid (n, x) \mapsto x_k u(n, x) \in \mathcal{Y}_{d,D}^i \text{ for all } k \in \{1, \dots, D\} \right\}. \end{aligned}$$

for functions on a mixed discrete-continuous state space. Here and below, \mathbb{R}_+^D denotes the set $[0, \infty)^D$.

3.3 Markov process and chemical master equation

The Markov process corresponding to the scaled and partitioned system is given by

$$\begin{pmatrix} X^{[M]}(t) \\ Y^{[M]}(t) \end{pmatrix} = \begin{pmatrix} n_0 \\ Mm_0 \end{pmatrix} + \sum_{j=1}^r \mathcal{P}_j \left(\int_0^t \alpha_j \left(X^{[M]}(s) \right) \beta_j^{[M]} \left(Y^{[M]}(s) \right) ds \right) \begin{pmatrix} \nu_j \\ \mu_j \end{pmatrix} \quad (24)$$

with $n_0 \in \mathbb{N}_0^d$ and $m_0 \in \mathbb{N}_0^D$. The probability distribution

$$p^{[M]}(t, n, m) = \mathbb{P} \left(X^{[M]}(t) = n, Y^{[M]}(t) = m \mid X^{[M]}(0) = n_0, Y^{[M]}(0) = Mm_0 \right)$$

is the solution of the CME

$$\begin{aligned} \partial_t p^{[M]}(t, n, m) &= \sum_{j=1}^r \left(\alpha_j(n - \nu_j) \beta_j^{[M]}(m - \mu_j) p^{[M]}(t, n - \nu_j, m - \mu_j) \right. \\ &\quad \left. - \alpha_j(n) \beta_j^{[M]}(m) p^{[M]}(t, n, m) \right) \end{aligned} \quad (25)$$

$$p^{[M]}(0, n, m) = \delta_{n_0}(n) \delta_{Mm_0}(m). \quad (26)$$

As before in (6), we let $p^{[M]}(t, n - \nu_j, m - \mu_j) = 0$ if $n - \nu_j \notin \mathbb{N}_0^d$ or $m - \mu_j \notin \mathbb{N}_0^D$. For the sake of a more compact notation we define the shift operators Δ_ν^1 and Δ_μ^2 with $\nu \in \mathbb{Z}^d$ and $\mu \in \mathbb{Z}^D$ via

$$\Delta_\nu^1 u(n, m) = \begin{cases} u(n - \nu, m) & \text{if } n - \nu \in \mathbb{N}_0^d \\ 0 & \text{else} \end{cases}, \quad (27)$$

$$\Delta_\mu^2 u(n, m) = \begin{cases} u(n, m - \mu) & \text{if } m - \mu \in \mathbb{N}_0^D \\ 0 & \text{else} \end{cases}. \quad (28)$$

The two shift operators commute, i.e.

$$\Delta_\nu^1 \Delta_\mu^2 u(n, m) = \Delta_\mu^2 \Delta_\nu^1 u(n, m) = \begin{cases} u(n - \nu, m - \mu) & \text{if } n - \nu \in \mathbb{N}_0^d, m - \mu \in \mathbb{N}_0^D \\ 0 & \text{else.} \end{cases}$$

We shall make the convention that applying a shift operator to a product $u(n, m)v(n, m)$ is to be understood as

$$(\Delta_\nu^1 uv)(n, m) = (\Delta_\nu^1(uv))(n, m) = u(n - \nu, m)v(n - \nu, m) = (\Delta_\nu^1 u)(\Delta_\nu^1 v)(n, m).$$

With these operators, the CME (25) can be reformulated as

$$\partial_t p^{[M]} = \sum_{j=1}^r (\Delta_{\nu_j}^1 \Delta_{\mu_j}^2 - 1) \alpha_j \beta_j^{[M]} p^{[M]}. \quad (29)$$

Lemma 2 *Let $\nu \in \mathbb{Z}^d$, $\mu \in \mathbb{Z}^D$, and let $u \in \ell_{d+D}^1$ be a function with the property that*

$$u(n, m) = 0 \quad \text{if } n + \nu \notin \mathbb{N}_0^d \text{ or } m + \mu \notin \mathbb{N}_0^D. \quad (30)$$

Then, the following assertions hold:

$$\begin{aligned}
(i) \quad & \sum_{n \in \mathbb{N}_0^d} ((\Delta_\nu^1 - 1)u)(n, m) = 0 \\
(ii) \quad & \sum_{m \in \mathbb{N}_0^D} ((\Delta_\mu^2 - 1)u)(n, m) = 0 \\
(iii) \quad & \sum_{n \in \mathbb{N}_0^d} \sum_{m \in \mathbb{N}_0^D} ((\Delta_\nu^1 \Delta_\mu^2 - 1)u)(n, m) = 0
\end{aligned}$$

If u is scalar-valued and $u \in \mathcal{X}_{d,D}^1$, then

$$(iv) \quad \sum_{m \in \mathbb{N}_0^D} m ((\Delta_\mu^2 - 1)u)(n, m) = \mu \sum_{m \in \mathbb{N}_0^D} u(n, m).$$

This lemma is not really new; similar considerations can be found, e.g., in [21, 15, 27]. However, since the lemma will be frequently used later on, we briefly recall the main argument of the proof.

Sketch of the Proof. For $\tilde{n} = n - \nu$ it follows from (27) that

$$\begin{aligned}
\sum_{n \in \mathbb{N}_0^d} ((\Delta_\nu^1 - 1)u)(n, m) &= \sum_{n \in \mathbb{N}_0^d} u(n - \nu, m) - \sum_{n \in \mathbb{N}_0^d} u(n, m) \\
&= \sum_{\tilde{n} + \nu \in \mathbb{N}_0^d} u(\tilde{n}, m) - \sum_{n \in \mathbb{N}_0^d} u(n, m) = 0
\end{aligned}$$

since according to (30) the two sums are identical. This proves (i), and (ii) follows by similar arguments. Then, (i) and (ii) imply (iii) via the identity

$$(\Delta_\nu^1 \Delta_\mu^2 - 1)u = \Delta_\nu^1 (\Delta_\mu^2 - 1)u + (\Delta_\nu^1 - 1)u.$$

Finally, (iv) is shown by substituting $\tilde{m} = m - \mu$:

$$\begin{aligned}
\sum_{m \in \mathbb{N}_0^D} m ((\Delta_\mu^2 - 1)u)(n, m) &= \sum_{m \in \mathbb{N}_0^D} mu(n, m - \mu) - \sum_{m \in \mathbb{N}_0^D} mu(n, m) \\
&= \sum_{\tilde{m} \in \mathbb{N}_0^D} (\tilde{m} + \mu)u(n, \tilde{m}) - \sum_{m \in \mathbb{N}_0^D} mu(n, m) \\
&= \mu \sum_{m \in \mathbb{N}_0^D} u(n, m). \quad \blacksquare
\end{aligned}$$

By Lemma (1), the function $\alpha_j \beta_j^{[M]} p^{[M]}$ satisfies (30). If $\alpha_j \beta_j^{[M]} p^{[M]} \in \ell_{d+D}^1$ we thus obtain that

$$\begin{aligned}
\sum_{n \in \mathbb{N}_0^d} \sum_{m \in \mathbb{N}_0^D} p^{[M]}(t, n, m) &= \sum_{n \in \mathbb{N}_0^d} \sum_{m \in \mathbb{N}_0^D} p^{[M]}(0, n, m) + \sum_{n \in \mathbb{N}_0^d} \sum_{m \in \mathbb{N}_0^D} \int_0^t \partial_t p^{[M]}(\tau, n, m) d\tau \\
&= 1 + \int_0^t \sum_{j=1}^r \sum_{n \in \mathbb{N}_0^d} \sum_{m \in \mathbb{N}_0^D} \left((\Delta_{\nu_j}^1 \Delta_{\mu_j}^2 - 1) \alpha_j \beta_j^{[M]} p^{[M]} \right) (\tau, n, m) = 1
\end{aligned}$$

Since it can be shown with standard arguments that $p(t, n, m) \geq 0$ (cf., e.g., Section 2.4 in [28]), this shows that $p^{[M]}(t, \cdot, \cdot)$ is a probability distribution at any time $t \geq 0$.

3.4 Piecewise deterministic process and Liouville-master equation

In order to formulate the piecewise deterministic process for general reaction systems, we define

$$\tilde{\beta}_j^{[M]}(x) = M^{1-\gamma(j)} \prod_{k=1}^D \frac{x_k^{\lambda_{jk}^{in}}}{\lambda_{jk}^{in}!}. \quad (31)$$

For $\gamma(j) = 1$, $\tilde{\beta}_j^{[M]}(x)$ is simply the term which appears on the right-hand side of the traditional reaction-rate equation (10). The following lemma shows that $\tilde{\beta}_j^{[M]}$ is closely related to $\beta_j^{[M]}$ defined in (23).

Lemma 3 *If $x \in \mathbb{R}_+^D$ and $x_k \geq 1/M$ for all $k \in \{1, \dots, D\}$, then there is a constant $C > 0$ such that*

$$\left| \beta_j^{[M]}(Mx) - \tilde{\beta}_j^{[M]}(x) \right| \leq CM^{-\gamma(j)}.$$

If $\lambda_{jk}^{in} \in \{0, 1\}$ for all $k = 1, \dots, D$, we even have $\beta_j^{[M]}(Mx) = \tilde{\beta}_j^{[M]}(x)$.

Proof. By Assumption 1 only the cases $\lambda_{jk}^{in} = 0, 1, 2$ must be considered. If $\lambda_{jk}^{in} \leq 1$, then

$$\left| M^{-\lambda_{jk}^{in}} \binom{Mx_k}{\lambda_{jk}^{in}} - \frac{x_k^{\lambda_{jk}^{in}}}{\lambda_{jk}^{in}!} \right| = 0$$

for any $j \in \{1, \dots, r\}$ and $k \in \{1, \dots, D\}$, such that the second assertion follows from (23) and (31). If $\lambda_{jk}^{in} = 2$ and $Mx_k > 1$, then

$$\left| M^{-\lambda_{jk}^{in}} \binom{Mx_k}{\lambda_{jk}^{in}} - \frac{x_k^{\lambda_{jk}^{in}}}{\lambda_{jk}^{in}!} \right| = \left| \frac{Mx_k(Mx_k - 1)}{2M^2} - \frac{x_k^2}{2} \right| = \frac{x_k}{2M},$$

whereas for $\lambda_{jk}^{in} = 2$ and $Mx_k < [0, 1]$ we obtain

$$\left| M^{-\lambda_{jk}^{in}} \binom{Mx_k}{\lambda_{jk}^{in}} - \frac{x_k^{\lambda_{jk}^{in}}}{\lambda_{jk}^{in}!} \right| = \frac{x_k^2}{2!} < \frac{1}{2M^2}.$$

Hence, for every $\lambda_{jk}^{in} \in \{0, 1, 2\}$ we have

$$\left| \binom{M^{-\lambda_{jk}^{in}} x_k}{\lambda_{jk}^{in}} - \frac{x_k^{\lambda_{jk}^{in}}}{\lambda_{jk}^{in}!} \right| \leq \frac{C}{2M},$$

and the assertion follows from

$$\left| \beta_j^{[M]}(Mx) - \tilde{\beta}_j^{[M]}(x) \right| = M^{1-\gamma(j)} \cdot \left| \prod_{k=1}^D M^{-\lambda_{jk}^{in}} \binom{Mx_k}{\lambda_{jk}^{in}} - \prod_{k=1}^D \frac{x_k^{\lambda_{jk}^{in}}}{\lambda_{jk}^{in!}} \right| \quad \blacksquare$$

Remark. According to Assumption 1, all $\beta_j^{[M]}$ and $\tilde{\beta}_j^{[M]}$ are polynomials of degree 2 or less. Hence, the Hessians $\nabla^2 \beta_j^{[M]}$ and $\nabla^2 \tilde{\beta}_j^{[M]}$ are both constant matrices, and higher derivatives vanish.

We are now ready to define the piecewise deterministic process corresponding to (24):

$$Z^{[M]}(t) = n_0 + \sum_{j=1}^r \mathcal{P}_j \left(\int_0^t \alpha_j \left(Z^{[M]}(s) \right) \tilde{\beta}_j^{[M]} \left(y^{[M]}(s) \right) ds \right) \nu_j \quad (32)$$

$$y^{[M]}(t) = m_0 + \frac{1}{M} \sum_{j=1}^r \left(\int_0^t \alpha_j \left(Z^{[M]}(s) \right) \tilde{\beta}_j^{[M]} \left(y^{[M]}(s) \right) ds \right) \mu_j \quad (33)$$

Example. For the reaction system defined in Table 3 with $n_0 = (10, 0)$ and $m_0 = 1$, the process (32)-(33) reduces to (15)-(16).

As before, the second equation (33) is piecewise deterministic in the sense that (33) is locally equivalent to the ODE

$$\dot{y}^{[M]}(t) = \frac{1}{M} \sum_{j=1}^r \alpha_j(z) \tilde{\beta}_j^{[M]} \left(y^{[M]}(t) \right) \mu_j$$

on time intervals where $Z^{[M]}(s) = z$ does not jump. The coupled equations (32)-(33) can be simulated, e.g., by adapting the Strang splitting method from Section 2.3.

The probability distribution associated to the original Markov process (24) evolves according to the CME (25) defined in the previous subsection. Now the question arises what the counterpart of the CME in case of the piecewise deterministic process (32)-(33) is. The answer can be found in [25, 8]: If

$$\int_S q^{[M]}(t, n, x) dx = \mathbb{P} \left(X^{[M]}(t) = n, y^{[M]}(t) \in S \mid X^{[M]}(0) = n_0, y^{[M]}(0) = m_0 \right)$$

for all measurable sets $S \subset \mathbb{R}_+^D$, then the density $q^{[M]}(t, n, x)$ is the solution of the Liouville-master equation (LME)

$$\begin{aligned} \partial_t q^{[M]}(t, n, x) &= \sum_{j \in J_1} \tilde{\beta}_j^{[M]}(x) \left(\alpha_j(n - \nu_j) q^{[M]}(t, n - \nu_j, x) - \alpha_j(n) q^{[M]}(t, n, x) \right) \\ &\quad - \frac{1}{M} \sum_{j=1}^r \alpha_j(n) \nabla \left(\tilde{\beta}_j^{[M]}(x) q^{[M]}(t, n, x) \right)^T \mu_j \end{aligned} \quad (34)$$

for $n \in \mathbb{N}_0^d$, $x \in \mathbb{R}_+^D$, $t \geq 0$, where ∇ acts on the x -variables only, i.e.

$$\nabla u(t, n, x) = \left(\frac{\partial u}{\partial x_1}(t, n, x), \dots, \frac{\partial u}{\partial x_D}(t, n, x) \right)^T.$$

The LME is provided with the boundary condition

$$q^{[M]}(t, n, x) = 0 \quad \text{if } x_i = 0 \text{ for one or more } i \in \{1, \dots, D\}. \quad (35)$$

and the initial data

$$q^{[M]}(0, n, x) = \delta_{n_0}(n)q_0(x). \quad (36)$$

Specifying $q_0(x)$ is somewhat delicate: since the process (32)-(33) starts in (n_0, m_0) with probability 1, $q_0(x)$ must be the corresponding Dirac delta, and then the question arises if a solution of (34)-(36) actually exists or in which sense the word “solution” has to be understood. Such analytical difficulties can be avoided if we choose $q_0 : \mathbb{R}_+^D \rightarrow \mathbb{R}$ to be a smooth nonnegative function with the properties

$$q_0(x) = 0 \quad \text{if } \|x - m_0\|_\infty > \varepsilon, \quad \int_{\mathbb{R}_+^D} q_0(x) = 1, \quad \int_{\mathbb{R}_+^D} xq_0(x) = m_0 \quad (37)$$

for some $0 < \varepsilon \ll 1$. If ε is sufficiently small, then the additional error caused by the modified initial data can be neglected.

With the shift operator defined in (27), the LME (34) can be reformulated as

$$\partial_t q^{[M]} = \sum_{j \in J_1} \tilde{\beta}_j^{[M]} (\Delta_{\nu_j}^1 - 1) \alpha_j q^{[M]} - \frac{1}{M} \sum_{j=1}^r \alpha_j \nabla \left(\tilde{\beta}_j^{[M]} q^{[M]} \right)^T \mu_j. \quad (38)$$

4 Error bound for the approximation by the piecewise deterministic process

In this section an error bound for the difference between the original Markov process and the piecewise deterministic process is presented. Convergence in distribution has recently been shown in [8]. In our analysis, only the *marginal* distribution of the discrete species $\mathcal{S}_1, \dots, \mathcal{S}_d$ and the conditional moments of the continuous species $\mathcal{S}_{d+1}, \dots, \mathcal{S}_D$ are considered, but in exchange, the *rate* of convergence will be established.

4.1 Marginal distributions, conditional expectations, and regularity assumptions

Assumption 2 We assume that the CME (29) with initial condition (26) has a unique classical solution $p^{[M]}(t, \cdot, \cdot) \in \mathcal{X}_{d,D}^3$ for $t \in [0, t_{end}]$ and that

$$(n, m) \mapsto \alpha_j(n)p^{[M]}(t, n, m) \in \mathcal{X}_{d,D}^3 \quad \text{for all } j \in \{1, \dots, r\}.$$

Moreover, we assume that the LME (38) with boundary condition (35) and initial data (36) has a unique classical solution $q^{[M]}(t, \cdot, \cdot) \in \mathcal{Y}_{d,D}^3$ for $t \in [0, t_{end}]$ and that

$$(n, x) \mapsto \alpha_j(n)q^{[M]}(t, n, x) \in \mathcal{Y}_{d,D}^3 \quad \text{for all } j \in \{1, \dots, r\}.$$

The marginal distribution of the discrete species $\mathcal{S}_1, \dots, \mathcal{S}_d$ is given by

$$p_1^{[M]}(t, n) = \sum_{m \in \mathbb{N}_0^D} p^{[M]}(t, n, m),$$

and we define

$$p_2^{[M]}(t, m|n) = \begin{cases} \frac{p^{[M]}(t, n, m)}{p_1^{[M]}(t, n)} & \text{if } p_1^{[M]}(t, n) > 0 \\ 0 & \text{else} \end{cases}.$$

If $p_1^{[M]}(t, n) > 0$, then $p_2^{[M]}(t, m|n)$ is the conditional probability that $Y^{[M]}(t) = m$ under the constraint that $X^{[M]}(t) = n$. Moreover, let

$$\eta^{[M]}(t, n) = \sum_{m \in \mathbb{N}_0^D} m p_2^{[M]}(t, m|n)$$

be the conditional expectation, and let

$$\sigma^{[M]}(t, n) = \sum_{m \in \mathbb{N}_0^D} \left(m - \eta^{[M]}(t, n) \right) \left(m - \eta^{[M]}(t, n) \right)^T p_2^{[M]}(t, m|n)$$

be the associated covariance matrix. The corresponding quantities for $q^{[M]}$ are defined as follows:

$$\begin{aligned} q_1^{[M]}(t, n) &= \int_{\mathbb{R}_+^D} q^{[M]}(t, n, x) dx \\ q_2^{[M]}(t, x|n) &= \begin{cases} \frac{q^{[M]}(t, n, x)}{q_1^{[M]}(t, n)} & \text{if } q_1^{[M]}(t, n) > 0 \\ 0 & \text{else} \end{cases} \\ \theta^{[M]}(t, n) &= \int_{\mathbb{R}_+^D} x q_2^{[M]}(t, x|n) dx \\ \varsigma(t, n) &= \int_{\mathbb{R}_+^D} \left(x - \theta^{[M]}(t, n) \right) \left(x - \theta^{[M]}(t, n) \right)^T q_2^{[M]}(t, x|n) dx \end{aligned}$$

Assumption 2 guarantees the existence of $\eta^{[M]}(t, n)$, $\sigma^{[M]}(t, n)$, $\theta^{[M]}(t, n)$, and $\varsigma(t, n)$ for all $n \in \mathbb{N}_0^d$ and $t \in [0, t_{\text{end}}]$

Assumption 3 *There is a constant $C > 0$ such that*

$$\|\eta^{[M]}(t, n)\| \leq C \cdot M, \quad \|\theta^{[M]}(t, n)\| \leq C \quad \text{for all } t \in [0, t_{\text{end}}], n \in \mathbb{N}_0^d \quad (39)$$

and

$$\|\sigma^{[M]}(t, n)\| \leq C \cdot M, \quad \|\varsigma^{[M]}(t, n)\| \leq \frac{C}{M} \quad \text{for all } t \in [0, t_{\text{end}}], n \in \mathbb{N}_0^d. \quad (40)$$

Moreover, all third central moments of $p_2^{[M]}(t, \cdot | n)$ are bounded by CM^2 , and all third central moments of $q_2^{[M]}(t, \cdot | n)$ are bounded by C/M for all $t \in [0, t_{\text{end}}]$ and $n \in \mathbb{N}_0^d$.

It follows from (26), (36), and (37) that $\eta^{[M]}(0|n) = \mathcal{O}(M)$, $\theta^{[M]}(0, n) = \mathcal{O}(1)$, and $\sigma^{[M]}(0|n) = 0$, and $\varsigma^{[M]}(0|n) = \mathcal{O}(\varepsilon^2)$. By a continuity argument, it can be shown that for every single M there is a $t_{\text{end}}^{[M]} > 0$ such that the above bounds hold for all $t \in [0, t_{\text{end}}^{[M]}]$. However, Assumption 3 is slightly stronger, because it is assumed that t_{end} does not depend on M .

At first sight, the assumptions on $\varsigma^{[M]}(t, n)$ and the third moments of $q_2^{[M]}(t, \cdot | n)$ seem to be somewhat restrictive. A closer inspection reveals, however, that the first term on the right-hand side of the LME (34) does not change these objects, and due to the factor M^{-1} of the second term, the above assumptions are reasonable.

Assumption 4 *Suppose that there is a constant $C > 0$ such that for all $t \in [0, t_{\text{end}}]$ and $j \in \{1, \dots, r\}$ the bound*

$$\|\alpha_j(\cdot)u(t, \cdot)\|_{\ell_a^1} \leq C \|u(t, \cdot)\|_{\ell_a^1} \quad (41)$$

holds for

$$\begin{aligned} u(t, n) &= \left(\beta_j^{[M]}(\eta^{[M]})p_1^{[M]} - \tilde{\beta}_j^{[M]}(\theta^{[M]})q_1^{[M]} \right) (t, n) \\ \text{or } u(t, n) &= \left(M^{-1}\beta_j^{[M]}(\eta^{[M]})\eta^{[M]}p_1^{[M]} - \tilde{\beta}_j^{[M]}(\theta^{[M]})\theta^{[M]}q_1^{[M]} \right) (t, n). \end{aligned}$$

Remark.

1. In both cases we have $\alpha_j(\cdot)u(t, \cdot) \in \ell_n^1$ according to Assumption 2.
2. Assumption 4 is true if $u(t, \cdot)$ decays sufficiently fast as $n \rightarrow \infty$. In the model problem discussed in Section 2.3 the assumption is true because there the state space is bounded in the n -direction.

4.2 A bound for the modeling error

After these preparations, the main result of this article can be formulated.

Theorem 1 *Let $p(t, n, m)$ be the solution of the CME (29) with initial condition (26), and let $q(t, n, x)$ be the solution of the LME (38) with boundary condition (35) and initial condition (36). Under the assumptions 1, 2, 3, and 4, there is a constant $C > 0$ such that the approximation error is bounded by*

$$\|p_1^{[M]}(t, \cdot) - q_1^{[M]}(t, \cdot)\|_{\ell_a^1} \leq \frac{C}{M} \quad (42)$$

$$\left\| \frac{1}{M}\eta^{[M]}(t, \cdot)p_1^{[M]}(t, \cdot) - \theta^{[M]}(t, \cdot)q_1^{[M]}(t, \cdot) \right\|_{\ell_a^1} \leq \frac{C}{M} \quad (43)$$

for all $t \in [0, t_{\text{end}}]$.

Remarks.

1. The choice of the parameter ε in (37) may alter the constants in the error bound and the value of t_{end} in Assumption 3. The convergence rate, however, is not affected by ε .
2. For the model problem from Section 2.3 the terms on the left-hand side of (42) and (43) have been estimated by numerical simulations for different values of M . The result was shown in Figure 2 and confirms that both errors are proportional to M^{-1} .

An immediate consequence of Theorem 1 is the following

Corollary 1 *If there is a constant $C_n^{\min} > 0$ such that*

$$p_1^{[M]}(t, n) \geq C_n^{\min} \quad \text{for all } t \in [0, t_{\text{end}}] \text{ and } M \in \mathbb{N},$$

then, under the assumptions of Theorem 1, the bound

$$\left\| \frac{1}{M} \eta^{[M]}(t, n) - \theta^{[M]}(t, n) \right\| \leq \frac{C}{C_n^{\min} M}$$

hold for the error in the conditional expectations.

Proof. Since

$$\frac{1}{M} \eta^{[M]} p_1^{[M]} - \theta^{[M]} q_1^{[M]} = \left(\frac{1}{M} \eta^{[M]} - \theta^{[M]} \right) p_1^{[M]} + \theta^{[M]} \left(p_1^{[M]} - q_1^{[M]} \right),$$

it follows from (43) that

$$\begin{aligned} & \left\| \frac{1}{M} \eta^{[M]}(t, n) - \theta^{[M]}(t, n) \right\| \\ &= \frac{1}{p_1^{[M]}(t, n)} \left(\left\| \frac{1}{M} \eta^{[M]}(t, n) p_1^{[M]}(t, n) - \theta^{[M]}(t, n) q_1^{[M]}(t, n) \right\| \right. \\ & \quad \left. + \|\theta^{[M]}(t, n)\| \cdot \left| p_1^{[M]}(t, n) - q_1^{[M]}(t, n) \right| \right) \\ & \leq \frac{C}{M p_1^{[M]}(t, n)}, \end{aligned}$$

and the assumption that $p_1^{[M]}(t, n) \geq C_n^{\min}$ yields the assertion. \blacksquare

For those n where $p_1^{[M]}(t, n) \approx 0$, the constant C_n^{\min} must be very small, and hence the difference between $\frac{1}{M} \eta^{[M]}(t, n)$ and $\theta^{[M]}(t, n)$ can be very large. However, this error is not relevant because $p_1^{[M]}(t, n) \approx 0$ means that these states are only visited with a very small probability.

Proof of Theorem 1. It will be shown in Lemma 5 and Lemma 6 that

$$E(t) = \|p_1^{[M]}(t, \cdot) - q_1^{[M]}(t, \cdot)\|_{\ell_a^1} + \left\| \frac{1}{M} \eta^{[M]}(t, \cdot) p_1^{[M]}(t, \cdot) - \theta^{[M]}(t, \cdot) q_1^{[M]}(t, \cdot) \right\|_{\ell_a^1}$$

satisfies the Gronwall inequality

$$E(t) \leq \frac{C_1}{M} + C_2 \int_0^t E(\tau) d\tau$$

for some constants $C_1, C_2 > 0$. Then, the Gronwall lemma yields

$$E(t) \leq \frac{C_1}{M} e^{C_2 t}$$

which implies (42) and (43). ■

Lemma 4 *Let $y : \mathbb{N}_0^d \longrightarrow \mathbb{R}^d$, $z : \mathbb{N}_0^d \longrightarrow \mathbb{R}^d$ with*

$$\max_{n \in \mathbb{N}_0^d} \|y(n)\| \leq CM, \quad \max_{n \in \mathbb{N}_0^d} \|z(n)\| \leq C, \quad (44)$$

and let $u \in \ell_d^1$ and $v \in \ell_d^1$. Then for every $j \in \{1, \dots, r\}$, there is a constant $C > 0$ such that

$$\left\| \beta_j^{[M]}(y)u - \tilde{\beta}_j^{[M]}(z)v \right\|_{\ell_d^1} \leq CM^{1-\gamma(j)} \left(\|M^{-1}yu - zv\|_{\ell_d^1} + \|u - v\|_{\ell_d^1} \right) + CM^{-\gamma(j)}$$

with $\gamma(j)$ defined in (22). Products of sequences are to be understood entry-wise, e.g.

$$\left[\beta_j^{[M]}(y)u \right] (n) = \beta_j^{[M]}(y(n))u(n),$$

and so on. Note that the assumption (44) implies that $M^{-1}yu - zv \in \ell_d^1$.

Proof. We investigate the propensities corresponding to the reaction scheme (1) and treat three cases separately:

- If $|\lambda_j^{in}|_1 = 0$, then $\beta_j^{[M]}(y) = \tilde{\beta}_j^{[M]}(y) = M^{1-\gamma(j)}$ is constant, and hence

$$\left\| \beta_j^{[M]}(y)u - \tilde{\beta}_j^{[M]}(z)v \right\|_{\ell_d^1} = M^{1-\gamma(j)} \|u - v\|_{\ell_d^1}.$$

- If $|\lambda_j^{in}|_1 = 1$, then there is a $k \in \{1, \dots, d\}$ such that $\beta_j^{[M]}(y) = M^{-\gamma(j)}y_k$ and $\tilde{\beta}_j^{[M]}(y) = M^{1-\gamma(j)}y_k$, and hence

$$\begin{aligned} \left\| \beta_j^{[M]}(y)u - \tilde{\beta}_j^{[M]}(z)v \right\| &= M^{1-\gamma(j)} \|M^{-1}y_k u - z_k v\|_{\ell_d^1} \\ &\leq M^{1-\gamma(j)} \|M^{-1}yu - zv\|_{\ell_d^1} \end{aligned}$$

- If $|\lambda_j^{in}|_1 = 2$, then the propensity $\tilde{\beta}_j^{[M]}(y)$ takes the form

$$\tilde{\beta}_j^{[M]}(y) = \hat{c}_j M^{1-\gamma(j)} y_k y_l \quad \text{with } \hat{c}_j = \begin{cases} c_j & \text{if } k \neq l \\ \frac{1}{2}c_j & \text{if } k = l \end{cases}$$

for some $k, l \in \{1, \dots, d\}$. The assumption (44) allows to apply Lemma 3, which gives

$$\left\| \beta_j^{[M]}(y)u - \tilde{\beta}_j^{[M]}(y/M)u \right\|_{\ell_d^1} \leq CM^{-\gamma(j)} \|u\|_{\ell_d^1}. \quad (45)$$

Thus, we only have to bound the difference

$$\begin{aligned} & \tilde{\beta}_j^{[M]}(y/M)u - \tilde{\beta}_j^{[M]}(z)v \\ &= \hat{c}_j M^{1-\gamma(j)} (M^{-2}y_k y_l u - z_k z_l v) \\ &= \hat{c}_j M^{1-\gamma(j)} \left(M^{-1}y_k (M^{-1}y_l u - z_l v) + z_l (M^{-1}y_k u - z_k v) - M^{-1}y_k z_l (u - v) \right). \end{aligned}$$

Since (44) implies that $|y_k(n)/M| \leq C$ and $|z_l(n)| \leq C$, it follows that

$$\left\| \tilde{\beta}_j^{[M]}(y/M)u - \tilde{\beta}_j^{[M]}(z)v \right\|_{\ell_d^1} \leq CM^{1-\gamma(j)} \left(\|M^{-1}yu - zv\|_{\ell_d^1} + \|u - v\|_{\ell_d^1} \right)$$

which, together with (45), proves the assertion. \blacksquare

Lemma 5 *Under the assumptions of Theorem 1 there is a constant $C > 0$ such that*

$$\begin{aligned} \|p_1^{[M]}(t, \cdot) - q_1^{[M]}(t, \cdot)\|_{\ell_d^1} &\leq \frac{C}{M} + C \int_0^t \left\| \left(\frac{1}{M} \eta^{[M]} p_1^{[M]} - \theta^{[M]} q_1^{[M]} \right) (\tau, \cdot) \right\|_{\ell_d^1} d\tau \\ &+ C \int_0^t \|p_1^{[M]}(\tau, \cdot) - q_1^{[M]}(\tau, \cdot)\|_{\ell_d^1} d\tau. \end{aligned}$$

for all $t \in [0, t_{end}]$.

Proof. Step 1: From the definition of the marginal distribution $p_1^{[M]}$ it follows that

$$\partial_t p_1^{[M]} = \sum_{j \in J_0} \sum_{m \in \mathbb{N}_0^D} (\Delta_{\nu_j}^1 \Delta_{\mu_j}^2 - 1) \alpha_j \beta_j^{[M]} p^{[M]} + \sum_{j \in J_1} \sum_{m \in \mathbb{N}_0^D} (\Delta_{\nu_j}^1 \Delta_{\mu_j}^2 - 1) \alpha_j \beta_j^{[M]} p^{[M]}.$$

The first sum vanishes, because $\Delta_{\nu_j}^1 = I$ for $j \in J_0$, and

$$\sum_{m \in \mathbb{N}_0^D} (\Delta_{\mu_j}^2 - 1) \alpha_j \beta_j^{[M]} p^{[M]} = 0$$

by Lemma 1 and Lemma 2. Since

$$\Delta_{\nu_j}^1 \Delta_{\mu_j}^2 - 1 = \Delta_{\nu_j}^1 (\Delta_{\mu_j}^2 - 1) + (\Delta_{\nu_j}^1 - 1)$$

and, again by Lemma 1 and Lemma 2,

$$\sum_{j \in J_1} \sum_{m \in \mathbb{N}_0^D} \Delta_{\nu_j}^1 (\Delta_{\mu_j}^2 - 1) \alpha_j \beta_j^{[M]} p^{[M]} = 0$$

we obtain

$$\begin{aligned}\partial_t p_1^{[M]} &= \sum_{j \in J_1} \sum_{m \in \mathbb{N}_0^D} (\Delta_{\nu_j}^1 - 1) \alpha_j \beta_j^{[M]} p^{[M]} \\ &= \sum_{j \in J_1} (\Delta_{\nu_j}^1 - 1) \alpha_j \sum_{m \in \mathbb{N}_0^D} \beta_j^{[M]} p_2^{[M]} p_1^{[M]}.\end{aligned}\quad (46)$$

Following the ideas of [14] we represent $\beta_j^{[M]}$ by the Taylor expansion

$$\begin{aligned}\beta_j^{[M]}(m) &= \beta_j^{[M]}(\eta^{[M]}) + \nabla \beta_j^{[M]}(\eta^{[M]})^T (m - \eta^{[M]}) \\ &\quad + \frac{1}{2} (m - \eta^{[M]})^T \left(\nabla^2 \beta_j^{[M]} \right) (m - \eta^{[M]})\end{aligned}\quad (47)$$

where $\eta^{[M]} = \eta^{[M]}(t, n)$. The usual remainder term vanishes because $\beta_j^{[M]}$ is at most quadratic due to Assumption 1. This yields

$$\sum_{m \in \mathbb{N}_0^D} \beta_j^{[M]}(m) p_2^{[M]}(t, m|n) = \beta_j^{[M]}(\eta^{[M]}) + \text{trace} \left(\sigma^{[M]}(t, n) \nabla^2 \beta_j^{[M]} \right) \quad (48)$$

because $\sum_{m \in \mathbb{N}_0^D} p_2^{[M]}(t, m|n) = 1$ and $\sum_{m \in \mathbb{N}_0^D} (m - \eta^{[M]}(t, n)) p_2^{[M]}(t, m|n) = 0$. According to (23) we have

$$\nabla^2 \beta_j^{[M]} = \begin{cases} 0 & \text{if } |\lambda_j^{in}|_1 < 2 \\ M^{-1-\gamma(j)} & \text{if } |\lambda_j^{in}|_1 = 2 \end{cases} \quad (49)$$

Since $\gamma(j) = 1$ for all $j \in J_1$ and $\sigma^{[M]} = \mathcal{O}(M)$ by assumption (40), it follows that $\sigma^{[M]} \nabla^2 \beta_j^{[M]} = \mathcal{O}(M^{-1})$ and thus (48) becomes

$$\sum_{m \in \mathbb{N}_0^D} \beta_j^{[M]}(m) p_2^{[M]}(t, m|n) = \beta_j^{[M]}(\eta^{[M]}) + \mathcal{O}(M^{-1}) \quad (50)$$

Substituting (50) into (46) gives

$$\partial_t p_1^{[M]} = \sum_{j \in J_1} (\Delta_{\nu_j}^1 - 1) \alpha_j \beta_j^{[M]}(\eta^{[M]}) p_1^{[M]} + \mathcal{O}(M^{-1}). \quad (51)$$

Step 2: Now the same steps are carried out *mutatis mutandis* for $q^{[M]}$. By definition of the marginal distribution $q_1^{[M]}$ and (38) we have

$$\begin{aligned}\partial_t q_1^{[M]} &= \sum_{j \in J_1} \int_{\mathbb{R}_+^D} (\Delta_{\nu_j}^1 - 1) \alpha_j \tilde{\beta}_j^{[M]} q^{[M]} dx - \frac{1}{M} \sum_{j=1}^r \alpha_j \int_{\mathbb{R}_+^D} \nabla \left(\tilde{\beta}_j^{[M]} q^{[M]} \right)^T \mu_j dx \\ &= \sum_{j \in J_1} (\Delta_{\nu_j}^1 - 1) \alpha_j \int_{\mathbb{R}_+^D} \tilde{\beta}_j^{[M]} q^{[M]} dx\end{aligned}$$

because the second term vanishes due to the boundary condition (36). Substituting the Taylor expansion

$$\begin{aligned} \tilde{\beta}_j^{[M]}(x) &= \tilde{\beta}_j^{[M]}(\theta^{[M]}) + \nabla \tilde{\beta}_j^{[M]}(\theta^{[M]})^T (x - \theta^{[M]}) \\ &\quad + \frac{1}{2} (x - \theta^{[M]})^T \left(\nabla^2 \tilde{\beta}_j^{[M]} \right) (x - \theta^{[M]}) \end{aligned} \quad (52)$$

and proceeding similar as before yields the counterpart of (48), namely

$$\int_{\mathbb{R}_+^D} \tilde{\beta}_j^{[M]}(x) q_2^{[M]}(t, x|n) dx = \tilde{\beta}_j^{[M]}(\theta^{[M]}) + \text{trace} \left(\varsigma^{[M]} \nabla^2 \tilde{\beta}_j^{[M]} \right). \quad (53)$$

Since $\varsigma^{[M]} \nabla^2 \tilde{\beta}_j^{[M]} = \mathcal{O}(M^{-1})$ by Assumption 3, it follows that

$$\partial_t q_1^{[M]} = \sum_{j \in J_1} (\Delta_{\nu_j}^1 - 1) \alpha_j \tilde{\beta}_j^{[M]}(\theta^{[M]}) q_1^{[M]} + \mathcal{O}(M^{-1}). \quad (54)$$

Step 3: Finally, we subtract (54) from (51) and integrate from 0 to t . Since $p_1^{[M]}(0) = q_1^{[M]}(0)$ and $\|(\Delta_{\nu_j}^1 - 1)\|_{\ell_d^1} \leq 2$, we obtain

$$\begin{aligned} &\|p_1^{[M]}(t, \cdot) - q_1^{[M]}(t, \cdot)\|_{\ell_d^1} \\ &\leq 2 \int_0^t \sum_{j \in J_1} \left\| \alpha_j \left(\beta_j^{[M]}(\eta^{[M]}) p_1^{[M]} - \tilde{\beta}_j^{[M]}(\theta^{[M]}) q_1^{[M]} \right) (\tau, \cdot) \right\|_{\ell_d^1} d\tau + \mathcal{O}(M^{-1}) \\ &\leq C \int_0^t \sum_{j \in J_1} \left\| \left(\beta_j^{[M]}(\eta^{[M]}) p_1^{[M]} - \tilde{\beta}_j^{[M]}(\theta^{[M]}) q_1^{[M]} \right) (\tau, \cdot) \right\|_{\ell_d^1} d\tau + \mathcal{O}(M^{-1}). \end{aligned}$$

The second inequality follows from Assumption 4. Now Lemma 4 can be applied, and since $M^{1-\gamma(j)} = 1$ for all $j \in J_1$, the assertion follows. \blacksquare

Lemma 6 *Under the assumptions of Theorem 1 there is a constant $C > 0$ such that*

$$\begin{aligned} \|M^{-1} \eta^{[M]} p_1^{[M]} - \theta^{[M]} q_1^{[M]}\|_{\ell_d^1} &\leq \frac{C}{M} + C \int_0^t \left\| \left[M^{-1} \eta^{[M]} p_1^{[M]} - \theta^{[M]} q_1^{[M]} \right] (\tau, \cdot) \right\|_{\ell_d^1} d\tau \\ &\quad + C \int_0^t \left\| \left[p_1^{[M]} - q_1^{[M]} \right] (\tau, \cdot) \right\|_{\ell_d^1} d\tau \end{aligned}$$

for all $t \in [0, t_{\text{end}}]$.

Proof. Step 1: From the definition of $\eta^{[M]}$ and $p_1^{[M]}$ and the CME (29) it follows that

$$\partial_t \left(\eta^{[M]} p_1^{[M]} \right) (t, n) = \sum_{m \in \mathbb{N}_0^D} m \partial_t p(t, n, m)$$

$$\begin{aligned}
&= \sum_{j=1}^r \sum_{m \in \mathbb{N}_0^D} m \left((\Delta_{\nu_j}^1 \Delta_{\mu_j}^2 - 1) \alpha_j \beta_j^{[M]} p^{[M]} \right) (t, n, m) \\
&= \sum_{j=1}^r \sum_{m \in \mathbb{N}_0^D} m \left(\Delta_{\nu_j}^1 (\Delta_{\mu_j}^2 - 1) \alpha_j \beta_j^{[M]} p^{[M]} \right) (t, n, m) \\
&\quad + \sum_{j=1}^r \sum_{m \in \mathbb{N}_0^D} m \left((\Delta_{\nu_j}^1 - 1) \alpha_j \beta_j^{[M]} p^{[M]} \right) (t, n, m) \\
&= \sum_{j=1}^r \mu_j \sum_{m \in \mathbb{N}_0^D} \left(\Delta_{\nu_j}^1 \alpha_j \beta_j^{[M]} p^{[M]} \right) (t, n, m) \tag{55} \\
&\quad + \sum_{j \in J_1} \sum_{m \in \mathbb{N}_0^D} m \left((\Delta_{\nu_j}^1 - 1) \alpha_j \beta_j^{[M]} p^{[M]} \right) (t, n, m) \tag{56}
\end{aligned}$$

due to Lemma 2 (iv) and the fact that $\Delta_{\nu_j}^1 - 1 = 0$ for all $j \in J_0$. For the first term (55), it follows from (48) that

$$\begin{aligned}
&\sum_{j=1}^r \mu_j \sum_{m \in \mathbb{N}_0^D} \left(\Delta_{\nu_j}^1 \alpha_j \beta_j^{[M]} p^{[M]} \right) (t, n, m) \\
&= \sum_{j=1}^r \mu_j \left(\Delta_{\nu_j}^1 \alpha_j \left[\beta_j^{[M]} (\eta^{[M]}) + \text{trace} \left(\sigma^{[M]} \nabla^2 \beta_j^{[M]} \right) \right] p_1^{[M]} \right) (t, n, m) \\
&= \sum_{j=1}^r \mu_j \left(\Delta_{\nu_j}^1 \alpha_j \beta_j^{[M]} (\eta^{[M]}) p_1^{[M]} \right) (t, n, m) + \mathcal{O}(1)
\end{aligned}$$

because (49) and Assumption (40) yield that

$$\left\| \text{trace} \left(\sigma^{[M]}(t, n) \nabla^2 \beta_j^{[M]} \right) \right\| \leq CM^{-\gamma(j)} \leq C$$

for all $t \in [0, t_{\text{end}}]$ and $n \in \mathbb{N}_0^d$. We remark that in the proof of Lemma 5, the same trace term only caused an error of $\mathcal{O}(M^{-1})$; cf. (50). This was due to the fact that in (51) we have $j \in J_1$ and hence $\gamma(j) = 1$, but this is not the case in (55)

For the second term (56), we use the Taylor expansion (47) to obtain

$$\sum_{m \in \mathbb{N}_0^D} m \beta_j^{[M]}(m) p^{[M]}(t, n, m) = \left(\beta_j^{[M]}(\eta^{[M]}) \eta^{[M]} + R_1(t, n) \right) p_1^{[M]}(t, n)$$

with $\eta^{[M]} = \eta^{[M]}(t, n)$ and the remainder term

$$\begin{aligned}
R_1(t, n) &= \sum_{m \in \mathbb{N}_0^D} m \left(\nabla \beta_j^{[M]}(\eta^{[M]})^T (m - \eta^{[M]}) \right. \\
&\quad \left. + \frac{1}{2} (m - \eta^{[M]})^T \left(\nabla^2 \beta_j^{[M]} \right) (m - \eta^{[M]}) \right) p_2^{[M]}(t, m|n)
\end{aligned}$$

$$\begin{aligned}
&= \sigma^{[M]} \nabla \beta_j^{[M]}(\eta^{[M]}) + \frac{1}{2} \eta^{[M]} \text{trace} \left(\sigma^{[M]} \nabla^2 \beta_j^{[M]} \right) \\
&\quad + \frac{1}{2} \sum_{m \in \mathbb{N}_0^D} (m - \eta^{[M]}) \left((m - \eta^{[M]})^T \left(\nabla^2 \beta_j^{[M]} \right) (m - \eta^{[M]}) \right) p_2^{[M]}(t, m|n)
\end{aligned}$$

which depends on the third moment of $p_2^{[M]}(t, \cdot|n)$. Now the crucial observation is that the sum in (56) is only taken over J_1 . For all $j \in J_1$, it follows from (23) and assumption (39) that

$$\nabla \beta_j^{[M]}(\eta^{[M]}) = \mathcal{O}(M^{-1}), \quad \nabla^2 \beta_j^{[M]} = \mathcal{O}(M^{-2}),$$

and Assumption 3 yields that $\|R_1(t, n)\| \leq C$ for all $n \in \mathbb{N}_0^d$ and $t \in [0, t_{\text{end}}]$. All in all, we have shown in step 1 that

$$\begin{aligned}
\partial_t \left(\eta^{[M]} p_1^{[M]} \right) (t, n) &= \sum_{j=1}^r \mu_j \left(\Delta_{\nu_j}^1 \alpha_j \beta_j^{[M]}(\eta^{[M]}) p_1^{[M]} \right) (t, n) \\
&\quad + \sum_{j \in J_1} \left((\Delta_{\nu_j}^1 - 1) \alpha_j \beta_j^{[M]}(\eta^{[M]}) \eta^{[M]} p_1^{[M]} \right) (t, n) + \mathcal{O}(1). \tag{57}
\end{aligned}$$

Later this equation will be multiplied by M^{-1} such that the rest term $\mathcal{O}(1)$ becomes $\mathcal{O}(M^{-1})$.

Step 2: Now an analogous equation is derived for $\partial_t \left(\theta^{[M]} q_1^{[M]} \right)$. According to the LME (38) we have

$$\begin{aligned}
\partial_t \left(\theta^{[M]} q_1^{[M]} \right) &= \int_{\mathbb{R}_+^D} x \partial_t q^{[M]} dx \\
&= \sum_{j \in J_1} \int_{\mathbb{R}_+^D} x \tilde{\beta}_j^{[M]} (\Delta_{\nu_j}^1 - 1) \alpha_j q^{[M]} dx - \frac{1}{M} \sum_{j=1}^r \int_{\mathbb{R}_+^D} x \alpha_j \nabla \left(\tilde{\beta}_j^{[M]} q^{[M]} \right)^T \mu_j dx
\end{aligned}$$

with $\tilde{\beta}_j^{[M]} = \tilde{\beta}_j^{[M]}(x)$ and $q^{[M]} = q^{[M]}(t, n, x)$. Integration by parts yields

$$\int_{\mathbb{R}_+^D} x \nabla \left(\tilde{\beta}_j^{[M]} q^{[M]} \right)^T \mu_j dx = -\mu_j \int_{\mathbb{R}_+^D} \tilde{\beta}_j^{[M]} q^{[M]} dx$$

because the boundary terms cancel due to the boundary condition (35) and $\lim_{x \rightarrow \infty} \tilde{\beta}_j^{[M]}(x) q^{[M]}(t, n, x) = 0$ (since $\tilde{\beta}_j^{[M]}(\cdot) q^{[M]}(t, n, \cdot) \in \ell_{d+D}^1$ due to Assumption 2). This gives

$$\partial_t \left(\theta^{[M]} q_1^{[M]} \right) = \sum_{j \in J_1} (\Delta_{\nu_j}^1 - 1) \alpha_j \int_{\mathbb{R}_+^D} x \tilde{\beta}_j^{[M]} q^{[M]} dx \tag{58}$$

$$+ \frac{1}{M} \sum_{j=1}^r \mu_j \int_{\mathbb{R}_+^D} \alpha_j \tilde{\beta}_j^{[M]} q^{[M]} dx. \tag{59}$$

Next, we insert the Taylor expansion (52) into (58) and obtain

$$\int_{\mathbb{R}_+^D} x \tilde{\beta}_j^{[M]} q^{[M]} dx = \left(\tilde{\beta}_j^{[M]}(\theta^{[M]}) \theta^{[M]} + R_2(t, n) \right) q_1^{[M]}(t, n)$$

with $\theta^{[M]} = \theta^{[M]}(t, n)$ and the rest term

$$\begin{aligned} R_2(t, n) &= \int_{\mathbb{R}_+^D} x \left(\nabla \tilde{\beta}_j^{[M]}(\theta^{[M]})^T (x - \theta^{[M]}) \right. \\ &\quad \left. + \frac{1}{2} (x - \theta^{[M]})^T \left(\nabla^2 \tilde{\beta}_j^{[M]} \right) (x - \theta^{[M]}) \right) q_2^{[M]} dx \\ &= \varsigma(t, n) \nabla \tilde{\beta}_j^{[M]}(\theta^{[M]}) + \frac{1}{2} \theta^{[M]} \text{trace} \left(\varsigma^{[M]}(t, n) \nabla^2 \tilde{\beta}_j^{[M]} \right) \\ &\quad + \frac{1}{2} \int_{\mathbb{R}_+^D} (x - \theta^{[M]}) (x - \theta^{[M]})^T \left(\nabla^2 \tilde{\beta}_j^{[M]} \right) (x - \theta^{[M]}) q_2^{[M]} dx. \end{aligned}$$

Again we note that the sum in (58) is only taken over J_1 , such that (31) and Assumption (39) imply that $\nabla \tilde{\beta}_j^{[M]}(\theta^{[M]})$ and its derivatives are uniformly bounded with respect to M . Combining this with Assumption 3 implies that $\|R_2(t, n)\| \leq C/M$ for all $n \in \mathbb{N}_0^d$ and $t \in [0, t_{\text{end}}]$.

Next, the second term (59) is investigated. The Taylor expansion (52) yields

$$\begin{aligned} &\frac{1}{M} \sum_{j=1}^r \mu_j \int_{\mathbb{R}_+^D} \alpha_j \tilde{\beta}_j^{[M]} q^{[M]} dx \\ &= \frac{1}{M} \sum_{j=1}^r \mu_j \alpha_j \left[\tilde{\beta}_j^{[M]}(\theta^{[M]}) + \text{trace} \left(\varsigma^{[M]} \nabla^2 \tilde{\beta}_j^{[M]} \right) \right] q_1^{[M]} \\ &= \frac{1}{M} \sum_{j=1}^r \mu_j \alpha_j \tilde{\beta}_j^{[M]}(\theta^{[M]}) q_1^{[M]} + \mathcal{O}(M^{-1}) \end{aligned}$$

because due to (31) and (40)

$$\left\| \text{trace} \left(\varsigma^{[M]}(t, n) \nabla^2 \tilde{\beta}_j^{[M]} \right) \right\| \leq C M^{-\gamma(j)} \leq C$$

for all $t \in [0, t_{\text{end}}]$ and $n \in \mathbb{N}_0^d$. We thus obtain

$$\begin{aligned} \partial_t \left(\theta^{[M]} q_1^{[M]} \right) &= \sum_{j \in J_1} (\Delta_{\nu_j}^1 - 1) \alpha_j \tilde{\beta}_j^{[M]}(\theta^{[M]}) \theta^{[M]} q_1^{[M]} \\ &\quad + \frac{1}{M} \sum_{j=1}^r \mu_j \left(\alpha_j \tilde{\beta}_j^{[M]}(\theta^{[M]}) q_1^{[M]} \right) (t, n, x) + \mathcal{O}(M^{-1}). \end{aligned} \tag{60}$$

Step 3: Multiplying (57) with M^{-1} , subtracting (60) and integrating from 0 to t yields, with $\|(\Delta_{\nu_j}^1 - 1)\|_{\ell_d^1} \leq 2$, that

$$\begin{aligned}
& \|M^{-1}\eta^{[M]}p_1^{[M]} - \theta^{[M]}q_1^{[M]}\|_{\ell_d^1} \\
& \leq \frac{C}{M} + \frac{C}{M} \int_0^t \sum_{j=1}^r \left\| \alpha_j \left[\beta_j^{[M]}(\eta^{[M]})p_1^{[M]} - \tilde{\beta}_j^{[M]}(\theta^{[M]})q_1^{[M]} \right] (\tau, \cdot) \right\|_{\ell_d^1} d\tau \\
& \quad + C \int_0^t \sum_{j \in J_1} \left\| \alpha_j \left[M^{-1}\beta_j^{[M]}(\eta^{[M]})\eta^{[M]}p_1^{[M]} - \tilde{\beta}_j^{[M]}(\theta^{[M]})\theta^{[M]}q_1^{[M]} \right] (\tau, \cdot) \right\|_{\ell_d^1} d\tau \\
& \leq \frac{C}{M} + \frac{C}{M} \int_0^t \sum_{j=1}^r \left\| \left[\beta_j^{[M]}(\eta^{[M]})p_1^{[M]} - \tilde{\beta}_j^{[M]}(\theta^{[M]})q_1^{[M]} \right] (\tau, \cdot) \right\|_{\ell_d^1} d\tau \\
& \quad + C \int_0^t \sum_{j \in J_1} \left\| \left[M^{-1}\beta_j^{[M]}(\eta^{[M]})\eta^{[M]}p_1^{[M]} - \tilde{\beta}_j^{[M]}(\theta^{[M]})\theta^{[M]}q_1^{[M]} \right] (\tau, \cdot) \right\|_{\ell_d^1} d\tau.
\end{aligned}$$

The second inequality follows from Assumption 4. Applying Lemma 4 yields

$$\begin{aligned}
& \frac{1}{M} \left\| \left[\beta_j^{[M]}(\eta^{[M]})p_1^{[M]} - \tilde{\beta}_j^{[M]}(\theta^{[M]})q_1^{[M]} \right] (\tau, \cdot) \right\|_{\ell_d^1} \\
& \leq C \left\| \left[M^{-1}\eta^{[M]}p_1^{[M]} - \theta^{[M]}q_1^{[M]} \right] (\tau, \cdot) \right\|_{\ell_d^1} + C \left\| \left[p_1^{[M]} - q_1^{[M]} \right] (\tau, \cdot) \right\|_{\ell_d^1} + \frac{C}{M}.
\end{aligned}$$

Hence, it remains to show that for all $j \in J_1$

$$\begin{aligned}
& \left\| \left[M^{-1}\beta_j^{[M]}(\eta^{[M]})\eta^{[M]}p_1^{[M]} - \tilde{\beta}_j^{[M]}(\theta^{[M]})\theta^{[M]}q_1^{[M]} \right] (\tau, \cdot) \right\|_{\ell_d^1} \\
& \leq \frac{C}{M} + C \left\| \left[M^{-1}\eta^{[M]}p_1^{[M]} - \theta^{[M]}q_1^{[M]} \right] (\tau, \cdot) \right\|_{\ell_d^1} + C \left\| p_1^{[M]}(\tau, \cdot) - q_1^{[M]}(\tau, \cdot) \right\|_{\ell_d^1} \tag{61}
\end{aligned}$$

for all $\tau \in [0, t_{\text{end}}]$. We decompose

$$\begin{aligned}
& M^{-1}\beta_j^{[M]}(\eta^{[M]})\eta^{[M]}p_1^{[M]} - \tilde{\beta}_j^{[M]}(\theta^{[M]})\theta^{[M]}q_1^{[M]} \\
& = \beta_j^{[M]}(\eta^{[M]}) \cdot \left[M^{-1}\eta^{[M]}p_1^{[M]} - \theta^{[M]}q_1^{[M]} \right] + \beta_j^{[M]}(\eta^{[M]})\theta^{[M]} \left[q_1^{[M]} - p_1^{[M]} \right] \\
& \quad + \theta^{[M]} \left[\beta_j^{[M]}(\eta^{[M]})p_1^{[M]} - \tilde{\beta}_j^{[M]}(\theta^{[M]})q_1^{[M]} \right],
\end{aligned}$$

take the ℓ^1 -norm on both sides, apply the triangle inequality and use that

$$\begin{aligned}
\max_{\tau \in [0, t_{\text{end}}]} \max_{n \in \mathbb{N}_0^d} \left| \beta_j^{[M]}(\eta^{[M]}(\tau, n)) \right| & \leq C \quad \text{for all } j \in J_1, \\
\max_{\tau \in [0, t_{\text{end}}]} \max_{n \in \mathbb{N}_0^d} \left| \theta^{[M]}(\tau, n) \right| & \leq C
\end{aligned}$$

due to Assumption (39). The last term can be bounded by applying Lemma 4 (with $\gamma(j) = 1$ since $j \in J_1$). This proves (61) and completes the proof of Lemma 6. \blacksquare

References

- [1] A. Alfonsi, E. Cancès, G. Turinici, B. D. Ventura, and W. Huisinga. Adaptive simulation of hybrid stochastic and deterministic models for biochemical systems. *ESAIM Proceeding*, 14:1–13, 2005.
- [2] D. F. Anderson and D. J. Higham. Multi-level Monte Carlo for continuous time Markov chains, with applications in biochemical kinetics. *SIAM Multiscale Modeling and Simulation*, 2012.
- [3] D. F. Anderson and T. G. Kurtz. *Continuous time Markov chain models for chemical reaction networks*, chapter 1 in Design and Analysis of Biomolecular Circuits: Engineering Approaches to Systems and Synthetic Biology, pages 3–42. Springer, 2011.
- [4] K. Ball, T. G. Kurtz, L. Popovic, and G. Rempala. Asymptotic analysis of multiscale approximations to reaction networks. *Ann. Appl. Probab.*, 16(4):1925–1961, 2006.
- [5] K. Burrage and T. Tian. Poisson Runge-Kutta methods for chemical reaction systems. In Y. L. W. Sun and T. Tang, editors, *Advances in Scientific Computing and Applications*, pages 82–96. Science Press, Beijing/New York, 2004.
- [6] K. Burrage, T. Tian, and P. Burrage. A multi-scaled approach for simulating chemical reaction systems. *Prog. Biophys. Mol. Biol.*, 85:217–234, 2004.
- [7] Y. Cao, D. T. Gillespie, and L. R. Petzold. The slow-scale stochastic simulation algorithm. *J. Chem. Phys.*, 122(1):014116, 2005.
- [8] A. Crudu, A. Debussche, A. Muller, and O. Radulescu. Convergence of stochastic gene networks to hybrid piecewise deterministic processes. *Annals of Applied Probab.*, to appear, 2011.
- [9] A. Crudu, A. Debussche, and O. Radulescu. Hybrid stochastic simplifications for multiscale gene networks. *BMC Systems Biology*, 3(89):1–25, 2009.
- [10] M. H. A. Davis. Piecewise-deterministic Markov processes: A general class of non-diffusion stochastic models. *Journal of the Royal Statistical Society, Series B (Methodological)*, 46(3):353388, 1984.
- [11] W. E, D. Liu, and E. Vanden-Eijnden. Nested stochastic simulation algorithm for chemical kinetic systems with disparate rates. *J. Chem. Phys.*, 123:194107, 2005.
- [12] W. E, D. Liu, and E. Vanden-Eijnden. Nested stochastic simulation algorithms for chemical kinetic systems with multiple time scales. *J. Comput. Phys.*, 221(1):158–180, 2007.
- [13] M. B. Elowitz, E. D. Siggia, P. S. Swain, and A. J. Levine. Stochastic gene expression in a single cell. *Science*, 297:1183–1186, 2002.
- [14] S. Engblom. Computing the moments of high dimensional solutions of the master equation. *Appl. Math. Comput.*, 180(2):498–515, 2006.

- [15] S. Engblom. Spectral approximation of solutions to the chemical master equation. *J. Comput. Appl. Math.*, 229(1):208–221, 2009.
- [16] S. N. Ethier and T. G. Kurtz. *Markov processes. Characterization and convergence*. Wiley Series in Probability and Statistics. Hoboken, NJ: John Wiley & Sons, 2005.
- [17] M. A. Gibson and J. Bruck. Efficient exact stochastic simulation of chemical systems with many species and many channels. *J. Phys. Chem. A*, 104:1876–1889, 2000.
- [18] D. Gillespie and L. Petzold. *System Modelling in Cellular Biology*, chapter 16. Numerical Simulation for Biochemical Kinetics, pages 331–354. MIT Press, 2006.
- [19] D. T. Gillespie. A general method for numerically simulating the stochastic time evolution of coupled chemical reactions. *J. Comput. Phys.*, 22:403–434, 1976.
- [20] D. T. Gillespie. A rigorous derivation of the chemical master equation. *Physica A*, 188:404–425, 1992.
- [21] D. T. Gillespie. The chemical Langevin equation. *J. Chem. Phys.*, 113:297–306, 2000.
- [22] D. T. Gillespie. Approximate accelerated stochastic simulation of chemically reacting systems. *J. Chem. Phys.*, 115(4):1716–1733, 2001.
- [23] J. Goutsias. Quasiequilibrium approximation of fast reaction kinetics in stochastic biochemical systems. *J. Chem. Phys.*, (122):184102, 2005.
- [24] E. L. Haseltine and J. B. Rawlings. Approximate simulation of coupled fast and slow reactions for stochastic chemical kinetics. *J. Chem. Phys.*, 117(15):6959–6969, 2002.
- [25] T. Henzinger, M. Mateescu, L. Mikeev, and V. Wolf. Hybrid numerical solution of the chemical master equation. In P. Quaglia, editor, *Proceedings of the 8th International Conference on Computational Methods in Systems Biology (CMSB10)*, pages 55–65. ACM, 2010.
- [26] D. J. Higham. Modeling and simulating chemical reactions. *SIAM Rev.*, 50(2):347–368, 2008.
- [27] T. Jahnke. An adaptive wavelet method for the chemical master equation. *SIAM J. Sci. Comput.*, 31(6):4373–4394, 2010.
- [28] T. Jahnke. On reduced models for the chemical master equation. *SIAM Multiscale Model. Simul.*, 9(4):1646–1676, 2011.
- [29] T. Jahnke and D. Altıntan. Efficient simulation of discrete stochastic reaction systems with a splitting method. *BIT*, 50(4):797–822, 2010.
- [30] T. G. Kurtz. The relationship between stochastic and deterministic models of chemical reactions. *J. Chem. Phys.*, 57:2976–2978, 1973.
- [31] T. G. Kurtz. *Approximation of population processes*. Number 36 in CBMS-NSF Reg. Conf. Ser. Appl. Math. SIAM Philadelphia, Pennsylvania, 1981.

- [32] T. Li. Analysis of explicit tau-leaping schemes for simulating chemically reacting systems. *Multiscale Model. Simul.*, 6(2):417–436, 2007.
- [33] H. H. McAdams and A. P. Arkin. Stochastic mechanisms in gene expression. *PNAS*, 94:814–819, 1997.
- [34] H. H. McAdams and A. P. Arkin. It’s a noisy business! Genetic regulation at the nanomolar scale. *Trends Genet.*, 15:65–69, 1999.
- [35] C. V. Rao and A. P. Arkin. Stochastic chemical kinetics and the quasi-steady-state assumption: Application to the Gillespie algorithm. *J. Chem. Phys.*, 118(11):4999–5010, 2003.
- [36] M. Rathinam, L. R. Petzold, Y. Cao, and D. T. Gillespie. Consistency and stability of tau-leaping schemes for chemical reaction systems. *Multiscale Model. Simul.*, 4(3):867–895, 2005.
- [37] M. Riedler. Almost sure convergence of numerical approximations for piecewise deterministic Markov processes. Technical report, Heriot-Watt University, 2010.
- [38] H. Salis and Y. Kaznessis. Accurate hybrid simulation of a system of coupled chemical or biochemical reactions. *J. Chem. Phys.*, 122, 2005.
- [39] R. Srivastava, L. You, J. Summers, and J. Yin. Stochastic vs. deterministic modeling of intracellular viral kinetics. *J. Theor. Biol.*, 218:309–321, 2002.
- [40] S. Zeiser, U. Franz, O. Wittich, and V. Liebscher. Simulation of genetic networks modelled by piecewise deterministic Markov processes. *IET Syst Biol.*, 2(3):113–135, 2008.

Preprint Series DFG-SPP 1324

<http://www.dfg-spp1324.de>

Reports

- [1] R. Ramlau, G. Teschke, and M. Zhariy. A Compressive Landweber Iteration for Solving Ill-Posed Inverse Problems. Preprint 1, DFG-SPP 1324, September 2008.
- [2] G. Plonka. The Easy Path Wavelet Transform: A New Adaptive Wavelet Transform for Sparse Representation of Two-dimensional Data. Preprint 2, DFG-SPP 1324, September 2008.
- [3] E. Novak and H. Woźniakowski. Optimal Order of Convergence and (In-) Tractability of Multivariate Approximation of Smooth Functions. Preprint 3, DFG-SPP 1324, October 2008.
- [4] M. Espig, L. Grasedyck, and W. Hackbusch. Black Box Low Tensor Rank Approximation Using Fibre-Crosses. Preprint 4, DFG-SPP 1324, October 2008.
- [5] T. Bonesky, S. Dahlke, P. Maass, and T. Raasch. Adaptive Wavelet Methods and Sparsity Reconstruction for Inverse Heat Conduction Problems. Preprint 5, DFG-SPP 1324, January 2009.
- [6] E. Novak and H. Woźniakowski. Approximation of Infinitely Differentiable Multivariate Functions Is Intractable. Preprint 6, DFG-SPP 1324, January 2009.
- [7] J. Ma and G. Plonka. A Review of Curvelets and Recent Applications. Preprint 7, DFG-SPP 1324, February 2009.
- [8] L. Denis, D. A. Lorenz, and D. Tiede. Greedy Solution of Ill-Posed Problems: Error Bounds and Exact Inversion. Preprint 8, DFG-SPP 1324, April 2009.
- [9] U. Friedrich. A Two Parameter Generalization of Lions' Nonoverlapping Domain Decomposition Method for Linear Elliptic PDEs. Preprint 9, DFG-SPP 1324, April 2009.
- [10] K. Bredies and D. A. Lorenz. Minimization of Non-smooth, Non-convex Functionals by Iterative Thresholding. Preprint 10, DFG-SPP 1324, April 2009.
- [11] K. Bredies and D. A. Lorenz. Regularization with Non-convex Separable Constraints. Preprint 11, DFG-SPP 1324, April 2009.

- [12] M. Döhler, S. Kunis, and D. Potts. Nonequispaced Hyperbolic Cross Fast Fourier Transform. Preprint 12, DFG-SPP 1324, April 2009.
- [13] C. Bender. Dual Pricing of Multi-Exercise Options under Volume Constraints. Preprint 13, DFG-SPP 1324, April 2009.
- [14] T. Müller-Gronbach and K. Ritter. Variable Subspace Sampling and Multi-level Algorithms. Preprint 14, DFG-SPP 1324, May 2009.
- [15] G. Plonka, S. Tenorth, and A. Iske. Optimally Sparse Image Representation by the Easy Path Wavelet Transform. Preprint 15, DFG-SPP 1324, May 2009.
- [16] S. Dahlke, E. Novak, and W. Sickel. Optimal Approximation of Elliptic Problems by Linear and Nonlinear Mappings IV: Errors in L_2 and Other Norms. Preprint 16, DFG-SPP 1324, June 2009.
- [17] B. Jin, T. Khan, P. Maass, and M. Pidcock. Function Spaces and Optimal Currents in Impedance Tomography. Preprint 17, DFG-SPP 1324, June 2009.
- [18] G. Plonka and J. Ma. Curvelet-Wavelet Regularized Split Bregman Iteration for Compressed Sensing. Preprint 18, DFG-SPP 1324, June 2009.
- [19] G. Teschke and C. Borries. Accelerated Projected Steepest Descent Method for Nonlinear Inverse Problems with Sparsity Constraints. Preprint 19, DFG-SPP 1324, July 2009.
- [20] L. Grasedyck. Hierarchical Singular Value Decomposition of Tensors. Preprint 20, DFG-SPP 1324, July 2009.
- [21] D. Rudolf. Error Bounds for Computing the Expectation by Markov Chain Monte Carlo. Preprint 21, DFG-SPP 1324, July 2009.
- [22] M. Hansen and W. Sickel. Best m-term Approximation and Lizorkin-Triebel Spaces. Preprint 22, DFG-SPP 1324, August 2009.
- [23] F.J. Hickernell, T. Müller-Gronbach, B. Niu, and K. Ritter. Multi-level Monte Carlo Algorithms for Infinite-dimensional Integration on \mathbb{R}^N . Preprint 23, DFG-SPP 1324, August 2009.
- [24] S. Dereich and F. Heidenreich. A Multilevel Monte Carlo Algorithm for Lévy Driven Stochastic Differential Equations. Preprint 24, DFG-SPP 1324, August 2009.
- [25] S. Dahlke, M. Fornasier, and T. Raasch. Multilevel Preconditioning for Adaptive Sparse Optimization. Preprint 25, DFG-SPP 1324, August 2009.

- [26] S. Dereich. Multilevel Monte Carlo Algorithms for Lévy-driven SDEs with Gaussian Correction. Preprint 26, DFG-SPP 1324, August 2009.
- [27] G. Plonka, S. Tenorth, and D. Roşca. A New Hybrid Method for Image Approximation using the Easy Path Wavelet Transform. Preprint 27, DFG-SPP 1324, October 2009.
- [28] O. Koch and C. Lubich. Dynamical Low-rank Approximation of Tensors. Preprint 28, DFG-SPP 1324, November 2009.
- [29] E. Faou, V. Gradinaru, and C. Lubich. Computing Semi-classical Quantum Dynamics with Hagedorn Wavepackets. Preprint 29, DFG-SPP 1324, November 2009.
- [30] D. Conte and C. Lubich. An Error Analysis of the Multi-configuration Time-dependent Hartree Method of Quantum Dynamics. Preprint 30, DFG-SPP 1324, November 2009.
- [31] C. E. Powell and E. Ullmann. Preconditioning Stochastic Galerkin Saddle Point Problems. Preprint 31, DFG-SPP 1324, November 2009.
- [32] O. G. Ernst and E. Ullmann. Stochastic Galerkin Matrices. Preprint 32, DFG-SPP 1324, November 2009.
- [33] F. Lindner and R. L. Schilling. Weak Order for the Discretization of the Stochastic Heat Equation Driven by Impulsive Noise. Preprint 33, DFG-SPP 1324, November 2009.
- [34] L. Kämmerer and S. Kunis. On the Stability of the Hyperbolic Cross Discrete Fourier Transform. Preprint 34, DFG-SPP 1324, December 2009.
- [35] P. Cerejeiras, M. Ferreira, U. Kähler, and G. Teschke. Inversion of the noisy Radon transform on $SO(3)$ by Gabor frames and sparse recovery principles. Preprint 35, DFG-SPP 1324, January 2010.
- [36] T. Jahnke and T. Udrescu. Solving Chemical Master Equations by Adaptive Wavelet Compression. Preprint 36, DFG-SPP 1324, January 2010.
- [37] P. Kittipoom, G. Kutyniok, and W.-Q. Lim. Irregular Shearlet Frames: Geometry and Approximation Properties. Preprint 37, DFG-SPP 1324, February 2010.
- [38] G. Kutyniok and W.-Q. Lim. Compactly Supported Shearlets are Optimally Sparse. Preprint 38, DFG-SPP 1324, February 2010.

- [39] M. Hansen and W. Sickel. Best m -Term Approximation and Tensor Products of Sobolev and Besov Spaces – the Case of Non-compact Embeddings. Preprint 39, DFG-SPP 1324, March 2010.
- [40] B. Niu, F.J. Hickernell, T. Müller-Gronbach, and K. Ritter. Deterministic Multi-level Algorithms for Infinite-dimensional Integration on $\mathbb{R}^{\mathbb{N}}$. Preprint 40, DFG-SPP 1324, March 2010.
- [41] P. Kittipoom, G. Kutyniok, and W.-Q. Lim. Construction of Compactly Supported Shearlet Frames. Preprint 41, DFG-SPP 1324, March 2010.
- [42] C. Bender and J. Steiner. Error Criteria for Numerical Solutions of Backward SDEs. Preprint 42, DFG-SPP 1324, April 2010.
- [43] L. Grasedyck. Polynomial Approximation in Hierarchical Tucker Format by Vector-Tensorization. Preprint 43, DFG-SPP 1324, April 2010.
- [44] M. Hansen und W. Sickel. Best m -Term Approximation and Sobolev-Besov Spaces of Dominating Mixed Smoothness - the Case of Compact Embeddings. Preprint 44, DFG-SPP 1324, April 2010.
- [45] P. Binev, W. Dahmen, and P. Lamby. Fast High-Dimensional Approximation with Sparse Occupancy Trees. Preprint 45, DFG-SPP 1324, May 2010.
- [46] J. Ballani and L. Grasedyck. A Projection Method to Solve Linear Systems in Tensor Format. Preprint 46, DFG-SPP 1324, May 2010.
- [47] P. Binev, A. Cohen, W. Dahmen, R. DeVore, G. Petrova, and P. Wojtaszczyk. Convergence Rates for Greedy Algorithms in Reduced Basis Methods. Preprint 47, DFG-SPP 1324, May 2010.
- [48] S. Kestler and K. Urban. Adaptive Wavelet Methods on Unbounded Domains. Preprint 48, DFG-SPP 1324, June 2010.
- [49] H. Yserentant. The Mixed Regularity of Electronic Wave Functions Multiplied by Explicit Correlation Factors. Preprint 49, DFG-SPP 1324, June 2010.
- [50] H. Yserentant. On the Complexity of the Electronic Schrödinger Equation. Preprint 50, DFG-SPP 1324, June 2010.
- [51] M. Guillemard and A. Iske. Curvature Analysis of Frequency Modulated Manifolds in Dimensionality Reduction. Preprint 51, DFG-SPP 1324, June 2010.
- [52] E. Herrholz and G. Teschke. Compressive Sensing Principles and Iterative Sparse Recovery for Inverse and Ill-Posed Problems. Preprint 52, DFG-SPP 1324, July 2010.

- [53] L. Kämmerer, S. Kunis, and D. Potts. Interpolation Lattices for Hyperbolic Cross Trigonometric Polynomials. Preprint 53, DFG-SPP 1324, July 2010.
- [54] G. Kutyniok and W.-Q Lim. Shearlets on Bounded Domains. Preprint 54, DFG-SPP 1324, July 2010.
- [55] A. Zeiser. Wavelet Approximation in Weighted Sobolev Spaces of Mixed Order with Applications to the Electronic Schrödinger Equation. Preprint 55, DFG-SPP 1324, July 2010.
- [56] G. Kutyniok, J. Lemvig, and W.-Q Lim. Compactly Supported Shearlets. Preprint 56, DFG-SPP 1324, July 2010.
- [57] A. Zeiser. On the Optimality of the Inexact Inverse Iteration Coupled with Adaptive Finite Element Methods. Preprint 57, DFG-SPP 1324, July 2010.
- [58] S. Jokar. Sparse Recovery and Kronecker Products. Preprint 58, DFG-SPP 1324, August 2010.
- [59] T. Aboiyar, E. H. Georgoulis, and A. Iske. Adaptive ADER Methods Using Kernel-Based Polyharmonic Spline WENO Reconstruction. Preprint 59, DFG-SPP 1324, August 2010.
- [60] O. G. Ernst, A. Mugler, H.-J. Starkloff, and E. Ullmann. On the Convergence of Generalized Polynomial Chaos Expansions. Preprint 60, DFG-SPP 1324, August 2010.
- [61] S. Holtz, T. Rohwedder, and R. Schneider. On Manifolds of Tensors of Fixed TT-Rank. Preprint 61, DFG-SPP 1324, September 2010.
- [62] J. Ballani, L. Grasedyck, and M. Kluge. Black Box Approximation of Tensors in Hierarchical Tucker Format. Preprint 62, DFG-SPP 1324, October 2010.
- [63] M. Hansen. On Tensor Products of Quasi-Banach Spaces. Preprint 63, DFG-SPP 1324, October 2010.
- [64] S. Dahlke, G. Steidl, and G. Teschke. Shearlet Coorbit Spaces: Compactly Supported Analyzing Shearlets, Traces and Embeddings. Preprint 64, DFG-SPP 1324, October 2010.
- [65] W. Hackbusch. Tensorisation of Vectors and their Efficient Convolution. Preprint 65, DFG-SPP 1324, November 2010.
- [66] P. A. Cioica, S. Dahlke, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Spatial Besov Regularity for Stochastic Partial Differential Equations on Lipschitz Domains. Preprint 66, DFG-SPP 1324, November 2010.

- [67] E. Novak and H. Woźniakowski. On the Power of Function Values for the Approximation Problem in Various Settings. Preprint 67, DFG-SPP 1324, November 2010.
- [68] A. Hinrichs, E. Novak, and H. Woźniakowski. The Curse of Dimensionality for Monotone and Convex Functions of Many Variables. Preprint 68, DFG-SPP 1324, November 2010.
- [69] G. Kutyniok and W.-Q. Lim. Image Separation Using Shearlets. Preprint 69, DFG-SPP 1324, November 2010.
- [70] B. Jin and P. Maass. An Analysis of Electrical Impedance Tomography with Applications to Tikhonov Regularization. Preprint 70, DFG-SPP 1324, December 2010.
- [71] S. Holtz, T. Rohwedder, and R. Schneider. The Alternating Linear Scheme for Tensor Optimisation in the TT Format. Preprint 71, DFG-SPP 1324, December 2010.
- [72] T. Müller-Gronbach and K. Ritter. A Local Refinement Strategy for Constructive Quantization of Scalar SDEs. Preprint 72, DFG-SPP 1324, December 2010.
- [73] T. Rohwedder and R. Schneider. An Analysis for the DIIS Acceleration Method used in Quantum Chemistry Calculations. Preprint 73, DFG-SPP 1324, December 2010.
- [74] C. Bender and J. Steiner. Least-Squares Monte Carlo for Backward SDEs. Preprint 74, DFG-SPP 1324, December 2010.
- [75] C. Bender. Primal and Dual Pricing of Multiple Exercise Options in Continuous Time. Preprint 75, DFG-SPP 1324, December 2010.
- [76] H. Harbrecht, M. Peters, and R. Schneider. On the Low-rank Approximation by the Pivoted Cholesky Decomposition. Preprint 76, DFG-SPP 1324, December 2010.
- [77] P. A. Cioica, S. Dahlke, N. Döhring, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Adaptive Wavelet Methods for Elliptic Stochastic Partial Differential Equations. Preprint 77, DFG-SPP 1324, January 2011.
- [78] G. Plonka, S. Tenorth, and A. Iske. Optimal Representation of Piecewise Hölder Smooth Bivariate Functions by the Easy Path Wavelet Transform. Preprint 78, DFG-SPP 1324, January 2011.

- [79] A. Mugler and H.-J. Starkloff. On Elliptic Partial Differential Equations with Random Coefficients. Preprint 79, DFG-SPP 1324, January 2011.
- [80] T. Müller-Gronbach, K. Ritter, and L. Yaroslavtseva. A Derandomization of the Euler Scheme for Scalar Stochastic Differential Equations. Preprint 80, DFG-SPP 1324, January 2011.
- [81] W. Dahmen, C. Huang, C. Schwab, and G. Welper. Adaptive Petrov-Galerkin methods for first order transport equations. Preprint 81, DFG-SPP 1324, January 2011.
- [82] K. Grella and C. Schwab. Sparse Tensor Spherical Harmonics Approximation in Radiative Transfer. Preprint 82, DFG-SPP 1324, January 2011.
- [83] D.A. Lorenz, S. Schiffler, and D. Trede. Beyond Convergence Rates: Exact Inversion With Tikhonov Regularization With Sparsity Constraints. Preprint 83, DFG-SPP 1324, January 2011.
- [84] S. Dereich, M. Scheutzow, and R. Schottstedt. Constructive quantization: Approximation by empirical measures. Preprint 84, DFG-SPP 1324, January 2011.
- [85] S. Dahlke and W. Sickel. On Besov Regularity of Solutions to Nonlinear Elliptic Partial Differential Equations. Preprint 85, DFG-SPP 1324, January 2011.
- [86] S. Dahlke, U. Friedrich, P. Maass, T. Raasch, and R.A. Ressel. An adaptive wavelet method for parameter identification problems in parabolic partial differential equations. Preprint 86, DFG-SPP 1324, January 2011.
- [87] A. Cohen, W. Dahmen, and G. Welper. Adaptivity and Variational Stabilization for Convection-Diffusion Equations. Preprint 87, DFG-SPP 1324, January 2011.
- [88] T. Jahnke. On Reduced Models for the Chemical Master Equation. Preprint 88, DFG-SPP 1324, January 2011.
- [89] P. Binev, W. Dahmen, R. DeVore, P. Lamby, D. Savu, and R. Sharpley. Compressed Sensing and Electron Microscopy. Preprint 89, DFG-SPP 1324, March 2011.
- [90] P. Binev, F. Blanco-Silva, D. Blom, W. Dahmen, P. Lamby, R. Sharpley, and T. Vogt. High Quality Image Formation by Nonlocal Means Applied to High-Angle Annular Dark Field Scanning Transmission Electron Microscopy (HAADF-STEM). Preprint 90, DFG-SPP 1324, March 2011.
- [91] R. A. Ressel. A Parameter Identification Problem for a Nonlinear Parabolic Differential Equation. Preprint 91, DFG-SPP 1324, May 2011.

- [92] G. Kutyniok. Data Separation by Sparse Representations. Preprint 92, DFG-SPP 1324, May 2011.
- [93] M. A. Davenport, M. F. Duarte, Y. C. Eldar, and G. Kutyniok. Introduction to Compressed Sensing. Preprint 93, DFG-SPP 1324, May 2011.
- [94] H.-C. Kreuzler and H. Yserentant. The Mixed Regularity of Electronic Wave Functions in Fractional Order and Weighted Sobolev Spaces. Preprint 94, DFG-SPP 1324, June 2011.
- [95] E. Ullmann, H. C. Elman, and O. G. Ernst. Efficient Iterative Solvers for Stochastic Galerkin Discretizations of Log-Transformed Random Diffusion Problems. Preprint 95, DFG-SPP 1324, June 2011.
- [96] S. Kunis and I. Melzer. On the Butterfly Sparse Fourier Transform. Preprint 96, DFG-SPP 1324, June 2011.
- [97] T. Rohwedder. The Continuous Coupled Cluster Formulation for the Electronic Schrödinger Equation. Preprint 97, DFG-SPP 1324, June 2011.
- [98] T. Rohwedder and R. Schneider. Error Estimates for the Coupled Cluster Method. Preprint 98, DFG-SPP 1324, June 2011.
- [99] P. A. Cioica and S. Dahlke. Spatial Besov Regularity for Semilinear Stochastic Partial Differential Equations on Bounded Lipschitz Domains. Preprint 99, DFG-SPP 1324, July 2011.
- [100] L. Grasedyck and W. Hackbusch. An Introduction to Hierarchical (H-) Rank and TT-Rank of Tensors with Examples. Preprint 100, DFG-SPP 1324, August 2011.
- [101] N. Chegini, S. Dahlke, U. Friedrich, and R. Stevenson. Piecewise Tensor Product Wavelet Bases by Extensions and Approximation Rates. Preprint 101, DFG-SPP 1324, September 2011.
- [102] S. Dahlke, P. Oswald, and T. Raasch. A Note on Quarkonial Systems and Multi-level Partition of Unity Methods. Preprint 102, DFG-SPP 1324, September 2011.
- [103] A. Uschmajew. Local Convergence of the Alternating Least Squares Algorithm For Canonical Tensor Approximation. Preprint 103, DFG-SPP 1324, September 2011.
- [104] S. Kvaal. Multiconfigurational time-dependent Hartree method for describing particle loss due to absorbing boundary conditions. Preprint 104, DFG-SPP 1324, September 2011.

- [105] M. Guillemard and A. Iske. On Groupoid C^* -Algebras, Persistent Homology and Time-Frequency Analysis. Preprint 105, DFG-SPP 1324, September 2011.
- [106] A. Hinrichs, E. Novak, and H. Woźniakowski. Discontinuous information in the worst case and randomized settings. Preprint 106, DFG-SPP 1324, September 2011.
- [107] M. Espig, W. Hackbusch, A. Litvinenko, H. Matthies, and E. Zander. Efficient Analysis of High Dimensional Data in Tensor Formats. Preprint 107, DFG-SPP 1324, September 2011.
- [108] M. Espig, W. Hackbusch, S. Handschuh, and R. Schneider. Optimization Problems in Contracted Tensor Networks. Preprint 108, DFG-SPP 1324, October 2011.
- [109] S. Dereich, T. Müller-Gronbach, and K. Ritter. On the Complexity of Computing Quadrature Formulas for SDEs. Preprint 109, DFG-SPP 1324, October 2011.
- [110] D. Belomestny. Solving optimal stopping problems by empirical dual optimization and penalization. Preprint 110, DFG-SPP 1324, November 2011.
- [111] D. Belomestny and J. Schoenmakers. Multilevel dual approach for pricing American style derivatives. Preprint 111, DFG-SPP 1324, November 2011.
- [112] T. Rohwedder and A. Uschmajew. Local convergence of alternating schemes for optimization of convex problems in the TT format. Preprint 112, DFG-SPP 1324, December 2011.
- [113] T. Görner, R. Hielscher, and S. Kunis. Efficient and accurate computation of spherical mean values at scattered center points. Preprint 113, DFG-SPP 1324, December 2011.
- [114] Y. Dong, T. Görner, and S. Kunis. An iterative reconstruction scheme for photoacoustic imaging. Preprint 114, DFG-SPP 1324, December 2011.
- [115] L. Kämmerer. Reconstructing hyperbolic cross trigonometric polynomials by sampling along generated sets. Preprint 115, DFG-SPP 1324, February 2012.
- [116] H. Chen and R. Schneider. Numerical analysis of augmented plane waves methods for full-potential electronic structure calculations. Preprint 116, DFG-SPP 1324, February 2012.
- [117] J. Ma, G. Plonka, and M.Y. Hussaini. Compressive Video Sampling with Approximate Message Passing Decoding. Preprint 117, DFG-SPP 1324, February 2012.

- [118] D. Heinen and G. Plonka. Wavelet shrinkage on paths for scattered data denoising. Preprint 118, DFG-SPP 1324, February 2012.
- [119] T. Jahnke and M. Kreim. Error bound for piecewise deterministic processes modeling stochastic reaction systems. Preprint 119, DFG-SPP 1324, March 2012.