

DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

A short theory of the Rayleigh-Ritz method

Harry Yserentant

Preprint 142



Edited by

AG Numerik/Optimierung
Fachbereich 12 - Mathematik und Informatik
Philipps-Universität Marburg
Hans-Meerwein-Str.
35032 Marburg

DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

A short theory of the Rayleigh-Ritz method

Harry Yserentant

Preprint 142



The consecutive numbering of the publications is determined by their chronological order.

The aim of this preprint series is to make new research rapidly available for scientific discussion. Therefore, the responsibility for the contents is solely due to the authors. The publications will be distributed by the authors.

A short theory of the Rayleigh-Ritz method

Harry Yserentant

Dedicated to Rolf Dieter Grigorieff on occasion of his 75th birthday

Abstract — We present some new error estimates for the eigenvalues and eigenfunctions obtained by the Rayleigh-Ritz method, the common variational method to solve eigenproblems. The errors are bounded in terms of the error of the best approximation of the eigenfunction under consideration by functions in the ansatz space. In contrast to the classical theory, the approximation error of eigenfunctions other than the given one does not enter into these estimates. The estimates are based on a bound for the norm of a certain projection operator, e.g., in finite element methods for second order eigenvalue problems, the H^1 -norm of the L_2 -projection onto the finite element space.

2010 Mathematical subject classification: 65N25; 65N15; 65N30.

Keywords: Eigenvalues; Eigenfunctions; Error estimates.

1. Introduction

The Rayleigh-Ritz method is a variational method to solve the eigenvalue problem for elliptic differential operators, that is, to compute their eigenvalues and the corresponding eigenfunctions. It is the direct counterpart of the Ritz method for the solution of the assigned boundary value problems. The Rayleigh-Ritz method has the advantage of being based on minimal, very general assumptions and produces optimal solutions in terms of the approximation properties of the underlying trial spaces. The theory of the Rayleigh-Ritz method has to a large extent been developed in the context of finite element methods, see [1, 2, 14]. In a more recent paper [12], Knyazev and Osborn derived error estimates into which, in contrast to the older theory, only the best approximation error of the eigenfunction under consideration enters. Here we show that such estimates can be derived in a very simple way utilizing the stability of certain projection operators, in the case of second order problems, the H^1 -stability of the L_2 -projection onto the given ansatz space.

Error estimates of this type are important for adaptive methods since it is obviously much simpler to adapt a mesh to a single eigenfunction than to a whole invariant subspace. A second, more subtle reason is that the regularity of the eigenfunctions often considerably differs and that, for example, approximating the eigenfunction for the minimum eigenvalue can be much more difficult than for the second one. The most prominent example of this type is the eigenfunctions of the Hamilton operator of the hydrogen atom whose knowledge is basic for our understanding of chemistry. These eigenfunctions are classified by three numbers, the principal quantum number $n = 1, 2, \dots$, the angular momentum quantum number $\ell = 0, \dots, n - 1$, and the magnetic quantum number $m = -\ell, \dots, \ell$. The eigenvalues

Harry Yserentant

Institut für Mathematik, Technische Universität Berlin, 10623 Berlin, Germany

E-mail: yserentant@math.tu-berlin.de

$\lambda = -1/(2n^2)$ depend only on the principal quantum number. They are highly degenerate for larger n ; their multiplicity is n^2 . The regularity of the eigenfunctions is determined by the angular momentum quantum number ℓ . They are contained in the Sobolev spaces H^s for $s < \ell + 5/2$, but not for $s = \ell + 5/2$, see [15]. Similar phenomena occur near reentrant edges and corners. Consider, for example, the eigenvalue problem for the negative Laplace operator on the unit disk, centered at the origin, with a slit along the positive x -axis and Dirichlet boundary conditions on the circular line and the top of the slit and Neumann boundary conditions on its bottom. The eigenvalues and eigenfunctions are in this case characterized by integers $k = 0, 1, \dots$ and $\ell = 1, 2, \dots$ and read

$$\lambda_{k,\ell} = (\omega_{\alpha,\ell})^2, \quad u_{k,\ell}(r, \varphi) = \sin(\alpha\varphi) J_\alpha(\omega_{\alpha,\ell} r) \quad (1.1)$$

in polar coordinate representation. The functions J_α are the Bessel functions of first kind and fractional order $\alpha = (2k + 1)/4$. They behave near the origin like $\sim r^\alpha$. The constants $\omega_{\alpha,\ell}$ are the zeroes of these Bessel functions in increasing order. This problem has been used in [13] as test example for the adaptive solution of eigenproblems.

2. The Rayleigh-Ritz method: framework and assumptions

We start from the usual abstract framework with two real Hilbert spaces \mathcal{H}_0 and $\mathcal{H}_1 \subseteq \mathcal{H}_0$ and a symmetric, coercive, and bounded bilinear form $a : \mathcal{H}_1 \times \mathcal{H}_1 \rightarrow \mathbb{R}$. The inner product on \mathcal{H}_0 is denoted by (u, v) and the induced norm by $\|u\|_0$. We equip the space \mathcal{H}_1 for simplicity with the energy norm $\|u\|$ induced by the bilinear form $a(u, v)$. For convenience we assume that \mathcal{H}_1 is compactly embedded into \mathcal{H}_0 and that both spaces are infinite dimensional. Then there exists an infinite sequence $0 < \lambda_1 \leq \lambda_2 \leq \dots$ of eigenvalues of finite multiplicity tending to infinity and an assigned sequence of eigenvectors u_1, u_2, \dots in \mathcal{H}_1 for which

$$(u_k, u_\ell) = \delta_{k\ell}, \quad a(u_k, u_\ell) = \lambda_k \delta_{k\ell}. \quad (2.1)$$

The example that we have in mind are second order elliptic eigenvalue problems over bounded domains Ω . In this case, $\mathcal{H}_0 = L_2(\Omega)$, and \mathcal{H}_1 is a subspace of the Sobolev space $H^1(\Omega)$, depending on the boundary conditions. The central results of this paper are, however, of more general nature and also apply, for example, to eigenfunctions for eigenvalues below the essential spectrum. Such situations arise in quantum chemistry.

The aim is to approximate the eigenvalues λ_k and the vectors in the assigned eigenspaces. For this, one chooses an n -dimensional subspace \mathcal{S} of \mathcal{H}_1 . Then there exist discrete eigenvectors u'_1, u'_2, \dots, u'_n in \mathcal{S} for eigenvalues $0 < \lambda'_1 \leq \lambda'_2 \leq \dots \leq \lambda'_n$, satisfying the relations

$$(u'_k, u'_\ell) = \delta_{k\ell}, \quad a(u'_k, u'_\ell) = \lambda'_k \delta_{k\ell}. \quad (2.2)$$

As will be shown, the discrete eigenvalues λ'_k approximate then the original eigenvalues λ and the discrete eigenvectors u'_k the corresponding eigenvectors u in a sense explained later. This already fixes the method, which replicates the weak form

$$a(u, v) = \lambda (u, v), \quad v \in \mathcal{H}_1, \quad (2.3)$$

of the original eigenvalue problem and is determined by the choice of the subspace \mathcal{S} replacing its solution space \mathcal{H}_1 . Typical approximation spaces \mathcal{S} are finite element spaces.

We will measure the approximation properties of the chosen subspace \mathcal{S} in terms of the a -orthogonal projection operator $P : \mathcal{H}_1 \rightarrow \mathcal{S}$ defined by

$$a(Pu, v) = a(u, v), \quad v \in \mathcal{S}. \quad (2.4)$$

With respect to the energy norm the projection Pu is the best approximation of $u \in \mathcal{H}_1$ by an element of \mathcal{S} , which means that for all $v \in \mathcal{S}$

$$\|u - Pu\| \leq \|u - v\|. \quad (2.5)$$

Our main assumption is that the correspondingly defined \mathcal{H}_0 -orthogonal projection Q from \mathcal{H}_0 onto \mathcal{S} is stable in the energy norm, that is, that there exists a constant κ with

$$\|Qv\| \leq \kappa \|v\|, \quad v \in \mathcal{H}_1. \quad (2.6)$$

The constant κ must be independent of hidden discretization parameters. This does not follow from the general assumptions made above but holds in many important cases. Examples are spectral methods in which the approximation spaces \mathcal{S} are built up from eigenfunctions of a nearby eigenvalue problem, say, in the case of a second order problem, from eigenfunctions of the Laplace operator. In this case the \mathcal{H}_0 - respectively L_2 -orthogonal projection coincides with the H^1 -orthogonal projection and satisfies the condition by definition.

The finite element case is more complicated. For quasi-uniform meshes estimates of this type can be traced back to the seminal paper [3] of Bank and Dupont. Consider a second-order problem, let $\mathcal{H}_0 = L_2(\Omega)$, and assume that $\mathcal{H}_1 \subseteq H^1(\Omega)$. Let the functions in the finite element space satisfy an inverse inequality

$$|v|_{H^1} \lesssim h^{-1} \|v\|_{L_2}, \quad v \in \mathcal{S}. \quad (2.7)$$

Let $\Pi : \mathcal{H}_1 \rightarrow \mathcal{S}$ be a quasi-interpolation operator for which

$$\|u - \Pi u\|_{L_2} \lesssim h |u|_{H^1}, \quad |\Pi u|_{H^1} \lesssim |u|_{H^1} \quad (2.8)$$

holds for all functions u in the solution space \mathcal{H}_1 . Then

$$|Q(u - \Pi u)|_{H^1} \lesssim h^{-1} \|Q(u - \Pi u)\|_{L_2} \leq h^{-1} \|u - \Pi u\|_{L_2} \lesssim |u|_{H^1} \quad (2.9)$$

for all these functions u . Since $Q\Pi u = \Pi u$ this implies the desired estimate

$$|Qu|_{H^1} \lesssim |Q(u - \Pi u)|_{H^1} + |\Pi u|_{H^1} \lesssim |u|_{H^1}. \quad (2.10)$$

Unfortunately, the h in the inverse inequality (2.7) behaves like the minimum element diameter and the h in (2.8) like the maximum element diameter. If the ratio of the maximum and the minimum element diameter becomes large, the proof breaks down.

The H^1 -stability of the L_2 -projection can, however, actually be proven under much weaker conditions on the grids. Partly rather technical sufficient conditions for the H^1 -stability of the L_2 -projection are [7, 8, 9, 10]. In [11], the case of piecewise linear elements in two space dimensions is treated. The most comprehensive result of this type is due to Bank and Yserentant [5]. It is shown in this paper that the L_2 -projection remains H^1 -stable for example for the highly nonuniform grids that are generated by bisection-like refinement schemes like the red-green refinement in two [4] and three [6] space dimensions and Lagrangian type finite elements of polynomial order up to twelve in two and up to seven in three space dimensions.

For realistic meshes and gridsizes changing in the average, on larger scales less abruptly than in an extreme case possible with such refinement schemes, the argumentation from [5] works without any condition to the polynomial order of the finite elements or any further condition to the triangulation up to shape regularity. The results in [5] even cover the case of hp -like finite element methods as long as the maximum order of the finite elements remains limited. There is therefore less doubt that the L_2 -projection remains H^1 -stable in almost all cases of practical interest so that (2.6) is a reasonable assumption at least for finite element discretizations of second order elliptic eigenvalue problems.

3. The error estimates

We first turn to the approximation of the eigenfunctions or, in the abstract setting, the eigenvectors. Starting point of our considerations is the following representation of the error between an eigenvector $u \in \mathcal{H}_1$ for an eigenvalue λ and its best \mathcal{H}_0 -approximation by a linear combination of discrete eigenvectors u'_k for eigenvalues λ'_k in a given neighborhood Λ of λ .

Lemma 3.1. *The error between an eigenvector $u \in \mathcal{H}_1$ of the original problem for the eigenvalue λ and its \mathcal{H}_0 -orthogonal projection onto the space spanned by the discrete eigenvectors $u'_k \in \mathcal{S}$ for the eigenvalues λ'_k in a given neighborhood Λ of λ possesses the representation*

$$u - \sum_{\lambda'_k \in \Lambda} (u, u'_k) u'_k = R(u - Pu) + (I - Q)(u - Pu), \quad (3.1)$$

where the mapping $R : \mathcal{H}_0 \rightarrow \mathcal{S}$ is defined by the expression

$$Rf = \sum_{\lambda'_k \notin \Lambda} \frac{\lambda'_k}{\lambda'_k - \lambda} (f, u'_k) u'_k. \quad (3.2)$$

Proof. We first represent the expression on the left hand side of (3.1) in the form

$$u - \sum_{\lambda'_k \in \Lambda} (u, u'_k) u'_k = \sum_{\lambda'_k \notin \Lambda} (u, u'_k) u'_k + u - \sum_{k=1}^n (u, u'_k) u'_k,$$

and replace the inner products in the first sum on the right hand side by

$$(u, u'_k) = \frac{\lambda'_k}{\lambda'_k - \lambda} (u - Pu, u'_k).$$

This is possible as u is an eigenvector and the u'_k are discrete eigenvectors; therefore

$$\lambda (u, u'_k) = a(u, u'_k) = a(Pu, u'_k) = \lambda'_k (Pu, u'_k).$$

Since the \mathcal{H}_0 -orthogonal projection can be written as eigenvector expansion

$$Qf = \sum_{k=1}^n (f, u'_k) u'_k,$$

the error representation

$$u - \sum_{\lambda'_k \in \Lambda} (u, u'_k) u'_k = R(u - Pu) + (u - Qu)$$

follows. As $QPu = Pu$, this yields (3.1). □

A first consequence of the error representation (3.1) is the following error estimate in the \mathcal{H}_0 -norm that is still independent of the \mathcal{H}_1 -stability (2.6) of the \mathcal{H}_0 -orthogonal projection Q .

Theorem 3.1. *Let $u \in \mathcal{H}_1$ be an eigenvector for the eigenvalue λ and let Λ be an arbitrarily given neighborhood of this eigenvalue. Then*

$$\left\| u - \sum_{\lambda'_k \in \Lambda} (u, u'_k) u'_k \right\|_0 \leq \max(1, \gamma) \|u - Pu\|_0, \quad (3.3)$$

where $\gamma = 0$ if there is no discrete eigenvalue λ'_k outside Λ and γ is otherwise given by

$$\gamma = \max_{\lambda'_k \notin \Lambda} \left| \frac{\lambda'_k}{\lambda'_k - \lambda} \right|. \quad (3.4)$$

Proof. As $(u'_k, u'_\ell) = \delta_{k\ell}$, for all $f \in \mathcal{H}_0$

$$\|Rf\|_0^2 = \sum_{\lambda'_k \notin \Lambda} \left| \frac{\lambda'_k}{\lambda'_k - \lambda} (f, u'_k) \right|^2 \leq \gamma^2 \|Qf\|_0^2.$$

The proposition thus follows from the error representation (3.1) using the \mathcal{H}_0 -orthogonality of the terms $R(u - Pu)$ respectively $Q(u - Pu)$ and $(I - Q)(u - Pu)$. \square

Taking into account the energy norm or \mathcal{H}_1 -stability of the \mathcal{H}_0 -orthogonal projection Q , a very similar estimate of the primarily interesting energy norm error can be derived:

Theorem 3.2. *Let $u \in \mathcal{H}_1$ be an eigenvector for the eigenvalue λ and let Λ be an arbitrarily given neighborhood of this eigenvalue. Then*

$$\left\| u - \sum_{\lambda'_k \in \Lambda} (u, u'_k) u'_k \right\| \leq (1 + (\gamma + 1)\kappa) \|u - Pu\|, \quad (3.5)$$

where γ is the same constant as in the previous theorem and κ the constant from (2.6).

Proof. The proof is again based on the error representation (3.1) and transfers almost verbatim from that of the previous theorem. As $a(u'_k, u'_\ell) = \lambda'_k \delta_{k\ell}$, for all $f \in \mathcal{H}_0$

$$\|Rf\|^2 = \sum_{\lambda'_k \notin \Lambda} \lambda'_k \left| \frac{\lambda'_k}{\lambda'_k - \lambda} (f, u'_k) \right|^2 \leq \gamma^2 \|Qf\|^2.$$

The only difference is that one can no longer argue using the orthogonality of the single terms but has to switch to the triangle inequality. The bound (2.6) for the energy norm of the projection operator Q enters in form of the estimate

$$\|Q(u - Pu)\| \leq \kappa \|u - Pu\|$$

for the projection of the approximation error. \square

The larger the neighborhood Λ of the eigenvalue λ under consideration is chosen, the more discrete eigenvectors u'_k are used to approximate the assigned eigenvector u and the smaller the error is, but the less specific the relation between the original and the discrete eigenvectors becomes. To establish a correspondence between the continuous and discrete eigenvalues, we need a crude error estimate as the following one:

Lemma 3.2. *If \mathcal{E}_k , $k \leq n$, is the subspace of \mathcal{H}_1 spanned by the eigenvectors u_1, \dots, u_k of the original problem for the eigenvalues $\lambda_1 \leq \dots \leq \lambda_k$ and if there is an $\varepsilon < 1$ such that*

$$\|u - Pu\|_0 \leq \varepsilon \|u\|_0 \quad (3.6)$$

for all $u \in \mathcal{E}_k$, the following error estimate holds for the k -th eigenvalue:

$$\lambda_k \leq \lambda'_k \leq \frac{1}{(1 - \varepsilon)^2} \lambda_k \quad (3.7)$$

Proof. Let \mathcal{E}'_k be the k -dimensional subspace of \mathcal{S} that is spanned by the first k discrete eigenvectors u'_1, \dots, u'_k . By the min-max characterization of the eigenvalue λ_k then

$$\lambda_k \leq \max_{u \in \mathcal{E}'_k} \frac{a(u, u)}{(u, u)} = \lambda'_k.$$

The functions Pu_1, \dots, Pu_k are linearly independent of each other. If namely a linear combination of these functions vanishes, the corresponding linear combination of the functions u_1, \dots, u_k vanishes because of (3.6) too. By the min-max characterization of λ'_k therefore

$$\lambda'_k \leq \max_{u \in \mathcal{E}_k} \frac{a(Pu, Pu)}{(Pu, Pu)}.$$

As $a(Pu, Pu) \leq a(u, u)$ and $\|Pu\|_0 \geq (1 - \varepsilon)\|u\|_0$ for $u \in \mathcal{E}_k$, the upper estimate

$$\lambda'_k \leq \frac{1}{(1 - \varepsilon)^2} \max_{u \in \mathcal{E}_k} \frac{a(u, u)}{(u, u)} = \frac{1}{(1 - \varepsilon)^2} \lambda_k$$

for the discrete eigenvalue λ'_k follows. \square

The lemma ensures that the discrete eigenvalues λ'_k tend in the limit to their continuous counterparts λ_k . If the distance Δ of the eigenvalue λ under consideration to the neighboring eigenvalues of the original problem is sufficiently large, an adequate choice for the neighborhood Λ in (3.3), (3.5) is therefore the interval of length Δ with midpoint λ . Then

$$\gamma \leq \frac{2\lambda}{\Delta} + 1. \quad (3.8)$$

Asymptotically then only approximate eigenvectors u'_k for eigenvalues λ'_k tending to λ are taken into account. If the eigenvalue λ belongs to a cluster of closely neighboring eigenvalues, the neighborhood Λ should be chosen accordingly and (3.3) and (3.5) be interpreted as a result on the approximation by an element in the corresponding discrete invariant subspace. In any case, the approximation order in the given norms defined in a proper sense completely corresponds to that for a corresponding boundary value problem with solution u .

A similar result as for the eigenvectors holds for the eigenvalues:

Theorem 3.3. *Let $u \in \mathcal{H}_1$, $\|u\|_0 = 1$, be an eigenvector for the eigenvalue λ and assume that already a discrete eigenvalue $\lambda'_k \geq \lambda$ exists for which $\lambda'_k - \lambda \leq \lambda$. Then*

$$\min_{\lambda'_k \geq \lambda} (\lambda'_k - \lambda) \leq (1 + \alpha\kappa)^2 \|u - Pu\|^2, \quad (3.9)$$

where $\alpha = 1$ if there is no discrete eigenvalue $\lambda'_k < \lambda$ and α is otherwise given by

$$\alpha = \max_{\lambda'_k < \lambda} \frac{\lambda}{\lambda - \lambda'_k}. \quad (3.10)$$

Proof. Let u' be the best \mathcal{H}_0 -norm approximation

$$u' = \sum_{\lambda'_k \geq \lambda} (u, u'_k) u'_k$$

of u by a linear combination of the discrete eigenvectors u'_k for eigenvalues $\lambda'_k \geq \lambda$. Then

$$\|u - u'\|^2 = \lambda \|u - u'\|_0^2 + \sum_{\lambda'_k \geq \lambda} (\lambda'_k - \lambda) (u, u'_k)^2.$$

Since u' and $u - u'$ are by definition \mathcal{H}_0 -orthogonal, this implies

$$\|u - u'\|^2 \geq \min_{\lambda'_k \geq \lambda} (\lambda'_k - \lambda) \|u\|_0^2 + \left(\lambda - \min_{\lambda'_k \geq \lambda} (\lambda'_k - \lambda) \right) \|u - u'\|_0^2.$$

As the second term on the right hand side is by assumption nonnegative and $\|u\|_0 = 1$, thus

$$\min_{\lambda'_k \geq \lambda} (\lambda'_k - \lambda) \leq \|u - u'\|^2.$$

The assertion follows from this estimate analogously to the proof of Theorem 3.2, simply replacing the neighborhood Λ of the eigenvalue λ by the half-open interval $[\lambda, \infty)$. \square

For eigenvalues greater than the minimum eigenvalue, the size of the prefactors depends asymptotically on the separation of the eigenvalue under consideration from the smaller eigenvalues. The speed with which the discrete eigenvalues converge to their continuous counterparts is asymptotically determined by the speed with which the square of the best energy norm approximation error of the assigned eigenfunctions tends to zero. As with the estimates from Theorem 3.1 and Theorem 3.2, pollution effects arising from the approximation error for other eigenfunctions than the one under consideration do not occur.

References

- [1] I. Babuška and J. Osborn, *Finite element-Galerkin approximation of the eigenvalues and eigenvectors of selfadjoint problems*, Math. Comput., **52** (1989), pp. 275–297.
- [2] I. Babuška and J. Osborn, *Eigenvalue problems*, in: *Handbook of Numerical Analysis, Vol.II, Finite Element Methods (Part 1)* (P. Ciarlet and J. Lions, eds.), Elsevier, 1991, pp. 641–792.
- [3] R. E. Bank and T. Dupont, *An optimal order process for solving finite element equations*, Math. Comput., **36** (1981), pp. 35–51.
- [4] R. E. Bank, A. H. Sherman, and A. Weiser, *Refinement algorithms and data structures for regular local mesh refinement*, in: *Scientific Computing (Applications of Mathematics and Computing to the Physical Sciences)* (R. S. Stepleman, ed.), North Holland, 1983, pp. 3–17.
- [5] R. E. Bank and H. Yserentant, *On the H^1 -stability of the L_2 -projection onto finite element spaces*, Numer. Math., (2013), DOI 10.1007/s00211-013-0562-4.
- [6] J. Bey, *Tetrahedral grid refinement*, Computing, **55** (1995), pp. 355–378.
- [7] J. Bramble, J. Pasciak, and O. Steinbach, *On the stability of the L_2 -projection in $H^1(\Omega)$* , Math. Comp., **71** (2001), pp. 147–156.
- [8] C. Carstensen, *Merging the Bramble-Pasciak-Steinbach and the Crouzeix-Thomée criterion for H^1 -stability of the L_2 -projection onto finite element spaces*, Math. Comput., **71** (2001), pp. 157–163.

-
- [9] C. Carstensen, *An adaptive mesh-refining algorithm allowing for an H^1 -stable L_2 -projection onto Courant finite element spaces*, *Constr. Approx.*, **20** (2004), pp. 549–564.
- [10] M. Crouzeix and V. Thomée, *The stability in L_p and W_p^1 of the L_2 -projection onto finite element function spaces*, *Math. Comput.*, **48** (1987), pp. 521–532.
- [11] M. Karkulik, D. Pavlicek, and D. Praetorius, *On 2D newest vertex bisection: optimality of mesh closure and H^1 -stability of L_2 -projection*, *Constr. Approx.*, (2013), DOI 10.1007/s00365-013-9192-4.
- [12] A. Knyazev and J. Osborn, *New a priori FEM error estimates for eigenvalues*, *SIAM J. Numer. Anal.*, **43** (2006), pp. 2647–2667.
- [13] K. Neymeyr, *A posteriori error estimation for elliptic eigenproblems*, *Numer. Linear Algebra Appl.*, **9** (2002), pp. 263–279.
- [14] P. Raviart and J. Thomas, *Introduction à L'Analyse Numérique des Équations aux Dérivées Partielles*, Masson, Paris, 1983.
- [15] H. Yserentant, *The hyperbolic cross space approximation of electronic wavefunctions*, *Numer. Math.*, **105** (2007), pp. 659–690.

Appendix. Improved constants

The norm of any nontrivial bounded projection operator Q on a Hilbert space coincides with the norm of the complementary projection $I - Q$, a seemingly not so much known result. The crucial estimate (2.6) implies therefore the estimate

$$\|(I - Q)v\| \leq \kappa \|v\|, \quad v \in \mathcal{H}_1.$$

Before we discuss its implications, we give a short proof of the general, abstract result. This and many other proofs can be found in [Szyld, Numer. Algor. (2006):42, 309–323].

Theorem. *Let $Q : \mathcal{H} \rightarrow \mathcal{H}$ be a bounded, nontrivial projection operator on a Hilbert space \mathcal{H} , that is, a projection operator that is different from zero and the identity. Then*

$$\|I - Q\| = \|Q\|.$$

Moreover, the norms of both operators are greater than or equal one.

Proof. Let $Qu \neq 0$. As $\|Q(Qu)\| = \|Qu\|$ the existence of such an u implies $\|Q\| \geq 1$. Correspondingly $\|I - Q\| \geq 1$. Let $u \in \mathcal{H}$ now be an arbitrary unit vector and let $v = Qu$ and $w = u - Qu$. We first assume that both v and w are different from the zero vector. Let

$$\tilde{v} = \frac{\|w\|}{\|v\|} v, \quad \tilde{w} = \frac{\|v\|}{\|w\|} w.$$

Since $Q\tilde{v} = \tilde{v}$ and $(I - Q)\tilde{w} = \tilde{w}$, the norm of Qu can be written as

$$\|Qu\| = \|v\| = \|\tilde{w}\| = \|(I - Q)(\tilde{v} + \tilde{w})\|.$$

As the two factors in front of v and w cancel, the norm of the vector $\tilde{v} + \tilde{w}$ is given by

$$\|\tilde{v} + \tilde{w}\|^2 = \|w\|^2 + \|v\|^2 + 2 \operatorname{Re}(v, w) = \|v + w\|^2$$

As $v + w = u$ and $\|u\| = 1$, the vector $\tilde{v} + \tilde{w}$ is therefore a unit vector, too. Thus

$$\|Qu\| \leq \|I - Q\|.$$

This also holds if $v = 0$ or $w = 0$. In the first case, $Qu = 0$ and nothing has to be shown. In the second case, $Qu = u$ and therefore $\|Qu\| = 1$. As $\|I - Q\| \geq 1$, the estimate above thus also holds for this case and therefore for arbitrary unit vectors u . We conclude that

$$\|Q\| \leq \|I - Q\|.$$

The reverse direction is shown in the same way interchanging the role of Q and $I - Q$. \square

Using the above norm estimate, the estimate (3.5) from Theorem 3.2 can be improved to

$$\left\| u - \sum_{\lambda'_k \in \Lambda} (u, u'_k) u'_k \right\| \leq (\gamma + 1) \kappa \|u - Pu\|,$$

and the estimate (3.9) from Theorem 3.3 to

$$\min_{\lambda'_k \geq \lambda} (\lambda'_k - \lambda) \leq (\alpha \kappa)^2 \|u - Pu\|^2.$$

That is, the notorious term “1 +” no longer shows up.

Preprint Series DFG-SPP 1324

<http://www.dfg-spp1324.de>

Reports

- [1] R. Ramlau, G. Teschke, and M. Zhariy. A Compressive Landweber Iteration for Solving Ill-Posed Inverse Problems. Preprint 1, DFG-SPP 1324, September 2008.
- [2] G. Plonka. The Easy Path Wavelet Transform: A New Adaptive Wavelet Transform for Sparse Representation of Two-dimensional Data. Preprint 2, DFG-SPP 1324, September 2008.
- [3] E. Novak and H. Woźniakowski. Optimal Order of Convergence and (In-) Tractability of Multivariate Approximation of Smooth Functions. Preprint 3, DFG-SPP 1324, October 2008.
- [4] M. Espig, L. Grasedyck, and W. Hackbusch. Black Box Low Tensor Rank Approximation Using Fibre-Crosses. Preprint 4, DFG-SPP 1324, October 2008.
- [5] T. Bonesky, S. Dahlke, P. Maass, and T. Raasch. Adaptive Wavelet Methods and Sparsity Reconstruction for Inverse Heat Conduction Problems. Preprint 5, DFG-SPP 1324, January 2009.
- [6] E. Novak and H. Woźniakowski. Approximation of Infinitely Differentiable Multivariate Functions Is Intractable. Preprint 6, DFG-SPP 1324, January 2009.
- [7] J. Ma and G. Plonka. A Review of Curvelets and Recent Applications. Preprint 7, DFG-SPP 1324, February 2009.
- [8] L. Denis, D. A. Lorenz, and D. Tiede. Greedy Solution of Ill-Posed Problems: Error Bounds and Exact Inversion. Preprint 8, DFG-SPP 1324, April 2009.
- [9] U. Friedrich. A Two Parameter Generalization of Lions' Nonoverlapping Domain Decomposition Method for Linear Elliptic PDEs. Preprint 9, DFG-SPP 1324, April 2009.
- [10] K. Bredies and D. A. Lorenz. Minimization of Non-smooth, Non-convex Functionals by Iterative Thresholding. Preprint 10, DFG-SPP 1324, April 2009.
- [11] K. Bredies and D. A. Lorenz. Regularization with Non-convex Separable Constraints. Preprint 11, DFG-SPP 1324, April 2009.

- [12] M. Döhler, S. Kunis, and D. Potts. Nonequispaced Hyperbolic Cross Fast Fourier Transform. Preprint 12, DFG-SPP 1324, April 2009.
- [13] C. Bender. Dual Pricing of Multi-Exercise Options under Volume Constraints. Preprint 13, DFG-SPP 1324, April 2009.
- [14] T. Müller-Gronbach and K. Ritter. Variable Subspace Sampling and Multi-level Algorithms. Preprint 14, DFG-SPP 1324, May 2009.
- [15] G. Plonka, S. Tenorth, and A. Iske. Optimally Sparse Image Representation by the Easy Path Wavelet Transform. Preprint 15, DFG-SPP 1324, May 2009.
- [16] S. Dahlke, E. Novak, and W. Sickel. Optimal Approximation of Elliptic Problems by Linear and Nonlinear Mappings IV: Errors in L_2 and Other Norms. Preprint 16, DFG-SPP 1324, June 2009.
- [17] B. Jin, T. Khan, P. Maass, and M. Pidcock. Function Spaces and Optimal Currents in Impedance Tomography. Preprint 17, DFG-SPP 1324, June 2009.
- [18] G. Plonka and J. Ma. Curvelet-Wavelet Regularized Split Bregman Iteration for Compressed Sensing. Preprint 18, DFG-SPP 1324, June 2009.
- [19] G. Teschke and C. Borries. Accelerated Projected Steepest Descent Method for Nonlinear Inverse Problems with Sparsity Constraints. Preprint 19, DFG-SPP 1324, July 2009.
- [20] L. Grasedyck. Hierarchical Singular Value Decomposition of Tensors. Preprint 20, DFG-SPP 1324, July 2009.
- [21] D. Rudolf. Error Bounds for Computing the Expectation by Markov Chain Monte Carlo. Preprint 21, DFG-SPP 1324, July 2009.
- [22] M. Hansen and W. Sickel. Best m -term Approximation and Lizorkin-Triebel Spaces. Preprint 22, DFG-SPP 1324, August 2009.
- [23] F.J. Hickernell, T. Müller-Gronbach, B. Niu, and K. Ritter. Multi-level Monte Carlo Algorithms for Infinite-dimensional Integration on \mathbb{R}^N . Preprint 23, DFG-SPP 1324, August 2009.
- [24] S. Dereich and F. Heidenreich. A Multilevel Monte Carlo Algorithm for Lévy Driven Stochastic Differential Equations. Preprint 24, DFG-SPP 1324, August 2009.
- [25] S. Dahlke, M. Fornasier, and T. Raasch. Multilevel Preconditioning for Adaptive Sparse Optimization. Preprint 25, DFG-SPP 1324, August 2009.

- [26] S. Dereich. Multilevel Monte Carlo Algorithms for Lévy-driven SDEs with Gaussian Correction. Preprint 26, DFG-SPP 1324, August 2009.
- [27] G. Plonka, S. Tenorth, and D. Roşca. A New Hybrid Method for Image Approximation using the Easy Path Wavelet Transform. Preprint 27, DFG-SPP 1324, October 2009.
- [28] O. Koch and C. Lubich. Dynamical Low-rank Approximation of Tensors. Preprint 28, DFG-SPP 1324, November 2009.
- [29] E. Faou, V. Gradinaru, and C. Lubich. Computing Semi-classical Quantum Dynamics with Hagedorn Wavepackets. Preprint 29, DFG-SPP 1324, November 2009.
- [30] D. Conte and C. Lubich. An Error Analysis of the Multi-configuration Time-dependent Hartree Method of Quantum Dynamics. Preprint 30, DFG-SPP 1324, November 2009.
- [31] C. E. Powell and E. Ullmann. Preconditioning Stochastic Galerkin Saddle Point Problems. Preprint 31, DFG-SPP 1324, November 2009.
- [32] O. G. Ernst and E. Ullmann. Stochastic Galerkin Matrices. Preprint 32, DFG-SPP 1324, November 2009.
- [33] F. Lindner and R. L. Schilling. Weak Order for the Discretization of the Stochastic Heat Equation Driven by Impulsive Noise. Preprint 33, DFG-SPP 1324, November 2009.
- [34] L. Kämmerer and S. Kunis. On the Stability of the Hyperbolic Cross Discrete Fourier Transform. Preprint 34, DFG-SPP 1324, December 2009.
- [35] P. Cerejeiras, M. Ferreira, U. Kähler, and G. Teschke. Inversion of the noisy Radon transform on $SO(3)$ by Gabor frames and sparse recovery principles. Preprint 35, DFG-SPP 1324, January 2010.
- [36] T. Jahnke and T. Udrescu. Solving Chemical Master Equations by Adaptive Wavelet Compression. Preprint 36, DFG-SPP 1324, January 2010.
- [37] P. Kittipoom, G. Kutyniok, and W.-Q. Lim. Irregular Shearlet Frames: Geometry and Approximation Properties. Preprint 37, DFG-SPP 1324, February 2010.
- [38] G. Kutyniok and W.-Q. Lim. Compactly Supported Shearlets are Optimally Sparse. Preprint 38, DFG-SPP 1324, February 2010.

- [39] M. Hansen and W. Sickel. Best m -Term Approximation and Tensor Products of Sobolev and Besov Spaces – the Case of Non-compact Embeddings. Preprint 39, DFG-SPP 1324, March 2010.
- [40] B. Niu, F.J. Hickernell, T. Müller-Gronbach, and K. Ritter. Deterministic Multi-level Algorithms for Infinite-dimensional Integration on $\mathbb{R}^{\mathbb{N}}$. Preprint 40, DFG-SPP 1324, March 2010.
- [41] P. Kittipoom, G. Kutyniok, and W.-Q Lim. Construction of Compactly Supported Shearlet Frames. Preprint 41, DFG-SPP 1324, March 2010.
- [42] C. Bender and J. Steiner. Error Criteria for Numerical Solutions of Backward SDEs. Preprint 42, DFG-SPP 1324, April 2010.
- [43] L. Grasedyck. Polynomial Approximation in Hierarchical Tucker Format by Vector-Tensorization. Preprint 43, DFG-SPP 1324, April 2010.
- [44] M. Hansen und W. Sickel. Best m -Term Approximation and Sobolev-Besov Spaces of Dominating Mixed Smoothness - the Case of Compact Embeddings. Preprint 44, DFG-SPP 1324, April 2010.
- [45] P. Binev, W. Dahmen, and P. Lamby. Fast High-Dimensional Approximation with Sparse Occupancy Trees. Preprint 45, DFG-SPP 1324, May 2010.
- [46] J. Ballani and L. Grasedyck. A Projection Method to Solve Linear Systems in Tensor Format. Preprint 46, DFG-SPP 1324, May 2010.
- [47] P. Binev, A. Cohen, W. Dahmen, R. DeVore, G. Petrova, and P. Wojtaszczyk. Convergence Rates for Greedy Algorithms in Reduced Basis Methods. Preprint 47, DFG-SPP 1324, May 2010.
- [48] S. Kestler and K. Urban. Adaptive Wavelet Methods on Unbounded Domains. Preprint 48, DFG-SPP 1324, June 2010.
- [49] H. Yserentant. The Mixed Regularity of Electronic Wave Functions Multiplied by Explicit Correlation Factors. Preprint 49, DFG-SPP 1324, June 2010.
- [50] H. Yserentant. On the Complexity of the Electronic Schrödinger Equation. Preprint 50, DFG-SPP 1324, June 2010.
- [51] M. Guillemard and A. Iske. Curvature Analysis of Frequency Modulated Manifolds in Dimensionality Reduction. Preprint 51, DFG-SPP 1324, June 2010.
- [52] E. Herrholz and G. Teschke. Compressive Sensing Principles and Iterative Sparse Recovery for Inverse and Ill-Posed Problems. Preprint 52, DFG-SPP 1324, July 2010.

- [53] L. Kämmerer, S. Kunis, and D. Potts. Interpolation Lattices for Hyperbolic Cross Trigonometric Polynomials. Preprint 53, DFG-SPP 1324, July 2010.
- [54] G. Kutyniok and W.-Q Lim. Shearlets on Bounded Domains. Preprint 54, DFG-SPP 1324, July 2010.
- [55] A. Zeiser. Wavelet Approximation in Weighted Sobolev Spaces of Mixed Order with Applications to the Electronic Schrödinger Equation. Preprint 55, DFG-SPP 1324, July 2010.
- [56] G. Kutyniok, J. Lemvig, and W.-Q Lim. Compactly Supported Shearlets. Preprint 56, DFG-SPP 1324, July 2010.
- [57] A. Zeiser. On the Optimality of the Inexact Inverse Iteration Coupled with Adaptive Finite Element Methods. Preprint 57, DFG-SPP 1324, July 2010.
- [58] S. Jokar. Sparse Recovery and Kronecker Products. Preprint 58, DFG-SPP 1324, August 2010.
- [59] T. Aboiyar, E. H. Georgoulis, and A. Iske. Adaptive ADER Methods Using Kernel-Based Polyharmonic Spline WENO Reconstruction. Preprint 59, DFG-SPP 1324, August 2010.
- [60] O. G. Ernst, A. Mugler, H.-J. Starkloff, and E. Ullmann. On the Convergence of Generalized Polynomial Chaos Expansions. Preprint 60, DFG-SPP 1324, August 2010.
- [61] S. Holtz, T. Rohwedder, and R. Schneider. On Manifolds of Tensors of Fixed TT-Rank. Preprint 61, DFG-SPP 1324, September 2010.
- [62] J. Ballani, L. Grasedyck, and M. Kluge. Black Box Approximation of Tensors in Hierarchical Tucker Format. Preprint 62, DFG-SPP 1324, October 2010.
- [63] M. Hansen. On Tensor Products of Quasi-Banach Spaces. Preprint 63, DFG-SPP 1324, October 2010.
- [64] S. Dahlke, G. Steidl, and G. Teschke. Shearlet Coorbit Spaces: Compactly Supported Analyzing Shearlets, Traces and Embeddings. Preprint 64, DFG-SPP 1324, October 2010.
- [65] W. Hackbusch. Tensorisation of Vectors and their Efficient Convolution. Preprint 65, DFG-SPP 1324, November 2010.
- [66] P. A. Cioica, S. Dahlke, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Spatial Besov Regularity for Stochastic Partial Differential Equations on Lipschitz Domains. Preprint 66, DFG-SPP 1324, November 2010.

- [67] E. Novak and H. Woźniakowski. On the Power of Function Values for the Approximation Problem in Various Settings. Preprint 67, DFG-SPP 1324, November 2010.
- [68] A. Hinrichs, E. Novak, and H. Woźniakowski. The Curse of Dimensionality for Monotone and Convex Functions of Many Variables. Preprint 68, DFG-SPP 1324, November 2010.
- [69] G. Kutyniok and W.-Q. Lim. Image Separation Using Shearlets. Preprint 69, DFG-SPP 1324, November 2010.
- [70] B. Jin and P. Maass. An Analysis of Electrical Impedance Tomography with Applications to Tikhonov Regularization. Preprint 70, DFG-SPP 1324, December 2010.
- [71] S. Holtz, T. Rohwedder, and R. Schneider. The Alternating Linear Scheme for Tensor Optimisation in the TT Format. Preprint 71, DFG-SPP 1324, December 2010.
- [72] T. Müller-Gronbach and K. Ritter. A Local Refinement Strategy for Constructive Quantization of Scalar SDEs. Preprint 72, DFG-SPP 1324, December 2010.
- [73] T. Rohwedder and R. Schneider. An Analysis for the DIIS Acceleration Method used in Quantum Chemistry Calculations. Preprint 73, DFG-SPP 1324, December 2010.
- [74] C. Bender and J. Steiner. Least-Squares Monte Carlo for Backward SDEs. Preprint 74, DFG-SPP 1324, December 2010.
- [75] C. Bender. Primal and Dual Pricing of Multiple Exercise Options in Continuous Time. Preprint 75, DFG-SPP 1324, December 2010.
- [76] H. Harbrecht, M. Peters, and R. Schneider. On the Low-rank Approximation by the Pivoted Cholesky Decomposition. Preprint 76, DFG-SPP 1324, December 2010.
- [77] P. A. Cioica, S. Dahlke, N. Döhring, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Adaptive Wavelet Methods for Elliptic Stochastic Partial Differential Equations. Preprint 77, DFG-SPP 1324, January 2011.
- [78] G. Plonka, S. Tenorth, and A. Iske. Optimal Representation of Piecewise Hölder Smooth Bivariate Functions by the Easy Path Wavelet Transform. Preprint 78, DFG-SPP 1324, January 2011.

- [79] A. Mugler and H.-J. Starkloff. On Elliptic Partial Differential Equations with Random Coefficients. Preprint 79, DFG-SPP 1324, January 2011.
- [80] T. Müller-Gronbach, K. Ritter, and L. Yaroslavl'tseva. A Derandomization of the Euler Scheme for Scalar Stochastic Differential Equations. Preprint 80, DFG-SPP 1324, January 2011.
- [81] W. Dahmen, C. Huang, C. Schwab, and G. Welper. Adaptive Petrov-Galerkin methods for first order transport equations. Preprint 81, DFG-SPP 1324, January 2011.
- [82] K. Grella and C. Schwab. Sparse Tensor Spherical Harmonics Approximation in Radiative Transfer. Preprint 82, DFG-SPP 1324, January 2011.
- [83] D.A. Lorenz, S. Schiffler, and D. Trede. Beyond Convergence Rates: Exact Inversion With Tikhonov Regularization With Sparsity Constraints. Preprint 83, DFG-SPP 1324, January 2011.
- [84] S. Dereich, M. Scheutzow, and R. Schottstedt. Constructive quantization: Approximation by empirical measures. Preprint 84, DFG-SPP 1324, January 2011.
- [85] S. Dahlke and W. Sickel. On Besov Regularity of Solutions to Nonlinear Elliptic Partial Differential Equations. Preprint 85, DFG-SPP 1324, January 2011.
- [86] S. Dahlke, U. Friedrich, P. Maass, T. Raasch, and R.A. Ressel. An adaptive wavelet method for parameter identification problems in parabolic partial differential equations. Preprint 86, DFG-SPP 1324, January 2011.
- [87] A. Cohen, W. Dahmen, and G. Welper. Adaptivity and Variational Stabilization for Convection-Diffusion Equations. Preprint 87, DFG-SPP 1324, January 2011.
- [88] T. Jahnke. On Reduced Models for the Chemical Master Equation. Preprint 88, DFG-SPP 1324, January 2011.
- [89] P. Binev, W. Dahmen, R. DeVore, P. Lamby, D. Savu, and R. Sharpley. Compressed Sensing and Electron Microscopy. Preprint 89, DFG-SPP 1324, March 2011.
- [90] P. Binev, F. Blanco-Silva, D. Blom, W. Dahmen, P. Lamby, R. Sharpley, and T. Vogt. High Quality Image Formation by Nonlocal Means Applied to High-Angle Annular Dark Field Scanning Transmission Electron Microscopy (HAADF-STEM). Preprint 90, DFG-SPP 1324, March 2011.
- [91] R. A. Ressel. A Parameter Identification Problem for a Nonlinear Parabolic Differential Equation. Preprint 91, DFG-SPP 1324, May 2011.

- [92] G. Kutyniok. Data Separation by Sparse Representations. Preprint 92, DFG-SPP 1324, May 2011.
- [93] M. A. Davenport, M. F. Duarte, Y. C. Eldar, and G. Kutyniok. Introduction to Compressed Sensing. Preprint 93, DFG-SPP 1324, May 2011.
- [94] H.-C. Kreuzler and H. Yserentant. The Mixed Regularity of Electronic Wave Functions in Fractional Order and Weighted Sobolev Spaces. Preprint 94, DFG-SPP 1324, June 2011.
- [95] E. Ullmann, H. C. Elman, and O. G. Ernst. Efficient Iterative Solvers for Stochastic Galerkin Discretizations of Log-Transformed Random Diffusion Problems. Preprint 95, DFG-SPP 1324, June 2011.
- [96] S. Kunis and I. Melzer. On the Butterfly Sparse Fourier Transform. Preprint 96, DFG-SPP 1324, June 2011.
- [97] T. Rohwedder. The Continuous Coupled Cluster Formulation for the Electronic Schrödinger Equation. Preprint 97, DFG-SPP 1324, June 2011.
- [98] T. Rohwedder and R. Schneider. Error Estimates for the Coupled Cluster Method. Preprint 98, DFG-SPP 1324, June 2011.
- [99] P. A. Cioica and S. Dahlke. Spatial Besov Regularity for Semilinear Stochastic Partial Differential Equations on Bounded Lipschitz Domains. Preprint 99, DFG-SPP 1324, July 2011.
- [100] L. Grasedyck and W. Hackbusch. An Introduction to Hierarchical (H-) Rank and TT-Rank of Tensors with Examples. Preprint 100, DFG-SPP 1324, August 2011.
- [101] N. Chegini, S. Dahlke, U. Friedrich, and R. Stevenson. Piecewise Tensor Product Wavelet Bases by Extensions and Approximation Rates. Preprint 101, DFG-SPP 1324, September 2011.
- [102] S. Dahlke, P. Oswald, and T. Raasch. A Note on Quarkonial Systems and Multi-level Partition of Unity Methods. Preprint 102, DFG-SPP 1324, September 2011.
- [103] A. Uschmajew. Local Convergence of the Alternating Least Squares Algorithm For Canonical Tensor Approximation. Preprint 103, DFG-SPP 1324, September 2011.
- [104] S. Kvaal. Multiconfigurational time-dependent Hartree method for describing particle loss due to absorbing boundary conditions. Preprint 104, DFG-SPP 1324, September 2011.

- [105] M. Guillemard and A. Iske. On Groupoid C^* -Algebras, Persistent Homology and Time-Frequency Analysis. Preprint 105, DFG-SPP 1324, September 2011.
- [106] A. Hinrichs, E. Novak, and H. Woźniakowski. Discontinuous information in the worst case and randomized settings. Preprint 106, DFG-SPP 1324, September 2011.
- [107] M. Espig, W. Hackbusch, A. Litvinenko, H. Matthies, and E. Zander. Efficient Analysis of High Dimensional Data in Tensor Formats. Preprint 107, DFG-SPP 1324, September 2011.
- [108] M. Espig, W. Hackbusch, S. Handschuh, and R. Schneider. Optimization Problems in Contracted Tensor Networks. Preprint 108, DFG-SPP 1324, October 2011.
- [109] S. Dereich, T. Müller-Gronbach, and K. Ritter. On the Complexity of Computing Quadrature Formulas for SDEs. Preprint 109, DFG-SPP 1324, October 2011.
- [110] D. Belomestny. Solving optimal stopping problems by empirical dual optimization and penalization. Preprint 110, DFG-SPP 1324, November 2011.
- [111] D. Belomestny and J. Schoenmakers. Multilevel dual approach for pricing American style derivatives. Preprint 111, DFG-SPP 1324, November 2011.
- [112] T. Rohwedder and A. Uschmajew. Local convergence of alternating schemes for optimization of convex problems in the TT format. Preprint 112, DFG-SPP 1324, December 2011.
- [113] T. Görner, R. Hielscher, and S. Kunis. Efficient and accurate computation of spherical mean values at scattered center points. Preprint 113, DFG-SPP 1324, December 2011.
- [114] Y. Dong, T. Görner, and S. Kunis. An iterative reconstruction scheme for photoacoustic imaging. Preprint 114, DFG-SPP 1324, December 2011.
- [115] L. Kämmerer. Reconstructing hyperbolic cross trigonometric polynomials by sampling along generated sets. Preprint 115, DFG-SPP 1324, February 2012.
- [116] H. Chen and R. Schneider. Numerical analysis of augmented plane waves methods for full-potential electronic structure calculations. Preprint 116, DFG-SPP 1324, February 2012.
- [117] J. Ma, G. Plonka, and M.Y. Hussaini. Compressive Video Sampling with Approximate Message Passing Decoding. Preprint 117, DFG-SPP 1324, February 2012.

- [118] D. Heinen and G. Plonka. Wavelet shrinkage on paths for scattered data denoising. Preprint 118, DFG-SPP 1324, February 2012.
- [119] T. Jahnke and M. Kreim. Error bound for piecewise deterministic processes modeling stochastic reaction systems. Preprint 119, DFG-SPP 1324, March 2012.
- [120] C. Bender and J. Steiner. A-posteriori estimates for backward SDEs. Preprint 120, DFG-SPP 1324, April 2012.
- [121] M. Espig, W. Hackbusch, A. Litvinenkoy, H.G. Matthiesy, and P. Wähnert. Efficient low-rank approximation of the stochastic Galerkin matrix in tensor formats. Preprint 121, DFG-SPP 1324, May 2012.
- [122] O. Bokanowski, J. Garcke, M. Griebel, and I. Klompmaker. An adaptive sparse grid semi-Lagrangian scheme for first order Hamilton-Jacobi Bellman equations. Preprint 122, DFG-SPP 1324, June 2012.
- [123] A. Mugler and H.-J. Starkloff. On the convergence of the stochastic Galerkin method for random elliptic partial differential equations. Preprint 123, DFG-SPP 1324, June 2012.
- [124] P.A. Cioica, S. Dahlke, N. Döhring, U. Friedrich, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R.L. Schilling. On the convergence analysis of Rothe's method. Preprint 124, DFG-SPP 1324, July 2012.
- [125] P. Binev, A. Cohen, W. Dahmen, and R. DeVore. Classification Algorithms using Adaptive Partitioning. Preprint 125, DFG-SPP 1324, July 2012.
- [126] C. Lubich, T. Rohwedder, R. Schneider, and B. Vandereycken. Dynamical approximation of hierarchical Tucker and Tensor-Train tensors. Preprint 126, DFG-SPP 1324, July 2012.
- [127] M. Kovács, S. Larsson, and K. Urban. On Wavelet-Galerkin methods for semilinear parabolic equations with additive noise. Preprint 127, DFG-SPP 1324, August 2012.
- [128] M. Bachmayr, H. Chen, and R. Schneider. Numerical analysis of Gaussian approximations in quantum chemistry. Preprint 128, DFG-SPP 1324, August 2012.
- [129] D. Rudolf. Explicit error bounds for Markov chain Monte Carlo. Preprint 129, DFG-SPP 1324, August 2012.
- [130] P.A. Cioica, K.-H. Kim, K. Lee, and F. Lindner. On the $L_q(L_p)$ -regularity and Besov smoothness of stochastic parabolic equations on bounded Lipschitz domains. Preprint 130, DFG-SPP 1324, December 2012.

- [131] M. Hansen. n -term Approximation Rates and Besov Regularity for Elliptic PDEs on Polyhedral Domains. Preprint 131, DFG-SPP 1324, December 2012.
- [132] R. E. Bank and H. Yserentant. On the H^1 -stability of the L_2 -projection onto finite element spaces. Preprint 132, DFG-SPP 1324, December 2012.
- [133] M. Gnewuch, S. Mayer, and K. Ritter. On Weighted Hilbert Spaces and Integration of Functions of Infinitely Many Variables. Preprint 133, DFG-SPP 1324, December 2012.
- [134] D. Crisan, J. Diehl, P.K. Friz, and H. Oberhauser. Robust Filtering: Correlated Noise and Multidimensional Observation. Preprint 134, DFG-SPP 1324, January 2013.
- [135] Wolfgang Dahmen, Christian Plesken, and Gerrit Welper. Double Greedy Algorithms: Reduced Basis Methods for Transport Dominated Problems. Preprint 135, DFG-SPP 1324, February 2013.
- [136] Aicke Hinrichs, Erich Novak, Mario Ullrich, and Henryk Wozniakowski. The Curse of Dimensionality for Numerical Integration of Smooth Functions. Preprint 136, DFG-SPP 1324, February 2013.
- [137] Markus Bachmayr, Wolfgang Dahmen, Ronald DeVore, and Lars Grasedyck. Approximation of High-Dimensional Rank One Tensors. Preprint 137, DFG-SPP 1324, March 2013.
- [138] Markus Bachmayr and Wolfgang Dahmen. Adaptive Near-Optimal Rank Tensor Approximation for High-Dimensional Operator Equations. Preprint 138, DFG-SPP 1324, April 2013.
- [139] Felix Lindner. Singular Behavior of the Solution to the Stochastic Heat Equation on a Polygonal Domain. Preprint 139, DFG-SPP 1324, May 2013.
- [140] Stephan Dahlke, Dominik Lellek, Shiu Hong Lui, and Rob Stevenson. Adaptive Wavelet Schwarz Methods for the Navier-Stokes Equation. Preprint 140, DFG-SPP 1324, May 2013.
- [141] Jonas Ballani and Lars Grasedyck. Tree Adaptive Approximation in the Hierarchical Tensor Format. Preprint 141, DFG-SPP 1324, June 2013.
- [142] Harry Yserentant. A short theory of the Rayleigh-Ritz method. Preprint 142, DFG-SPP 1324, July 2013.